

## On the effect of Prandtl number to subgrid-scale heat flux models


F.X. Trias<sup>1</sup>, D. Santos<sup>1</sup>, J.A. Hopman<sup>1</sup>, A. Gorobets<sup>2</sup>, A. Oliva<sup>1</sup>

<sup>1</sup> Heat and Mass Transfer Technological Center, Technical University of Catalonia, C/Colom 11, 08222 Terrassa (Barcelona), francesc.xavier.trias@upc.edu

<sup>2</sup> Keldysh Institute of Applied Mathematics, 4A, Miusskaya Sq., Moscow 125047, Russia

**Keywords:** LES, SGS models, buoyancy-driven flows, turbulence, Prandtl number effect

In this work, we aim to shed light to the following research question: *can we find a subgrid-scale (SGS) heat flux model with good physical and numerical properties, such that we can obtain satisfactory predictions for buoyancy-driven turbulence at high Prandtl ( $Pr$ ) numbers?* This is motivated by (i) our findings showing the reasons for the lack of accuracy of existing SGS heat flux models for LES [1, 2] (see Figure 1, left) and (ii) the fact that the flow topology in the near-wall region changes dramatically at high- $Pr$  numbers [3]. Firstly, we plan to study *a priori* the capability of the models proposed in [2] to provide accurate approximations of the actual SGS heat flux both in the bulk and in the near-wall regions. To do so, direct numerical simulations (DNS) at high  $Pr$  will be carried out. Moreover, the availability of DNS results at  $Pr = 0.7$  (air) from our previous works [1, 2] (see Figure 1, right) will allow to study the near-wall  $Pr$ -number effects. In this regard, it may be possible that some additional corrections/modifications of the model will be needed. Secondly, we plan to study *a posteriori* the performance of these models. In this case, LES simulations will be carried out with the same code and results compared with the DNS data. In a first stage, we plan to study the (*a posteriori*) modelization effects in the bulk region. This can be done by properly refining in the near-wall region. Finally, *a posteriori* tests including the modelization effects of the near-wall region will be performed.



**Figure 1:** Left: alignment trends of the actual SGS heat flux,  $q$ . For details the reader is referred to our work [1]. Right: DNS of the air-filled RBC at  $Ra = 10^{10}$  studied in Refs. [1, 2].

## References

- [1] F. Dabbagh, F. X. Trias, A. Gorobets, and A. Oliva. A priori study of subgrid-scale features in turbulent Rayleigh-Bénard convection. *Physics of Fluids*, 29:105103, 2017.
- [2] F.X. Trias, F.Dabbagh, A.Gorobets, and C.Oliet. On a proper tensor-diffusivity model for large-eddy simulation of buoyancy-driven turbulence. *Flow, Turbulence and Combustion*, 105:393–414, 2020.
- [3] S. Yigit, J. Hasslberger, M. Klein, and N. Chakraborty. Near wall Prandtl number effects on velocity gradient invariants and flow topologies in turbulent Rayleigh-Bénard convection. *Nature Scientific Reports*, 10(1):14887, 2020.