

# Advancing Fractional Calculus: A Fixed-Point Strategy for Nonlinear Caputo-Hadamard Equations

Muath Awadalla<sup>1</sup> and Manigandan Murugesan<sup>2,3,\*</sup>

<sup>1</sup> Department of Mathematics and Statistics, College of Science, King Faisal University, Hofuf, Al Ahsa, Saudi Arabia

<sup>2</sup> Center for Nonlinear and Complex Networks, SRM TRP Engineering College, Tiruchirapalli, Tamil Nadu, India

<sup>3</sup> Center for Research, Easwari Engineering College, Chennai, Tamil Nadu, India

## INFORMATION

### Keywords:

Caputo-Hadamard fractional derivative  
Krasnoselskii fixed-point theorem  
Banach fixed-point theorem  
existence and uniqueness  
Ulam-Hyers stability  
fractional differential systems

### MSC:

34B15, 34B18, 34B37, 58E30.

DOI: 10.23967/j.rimni.2026.10.79081

Revista Internacional  
Métodos numéricos  
para cálculo y diseño en ingeniería

RIMNI



UNIVERSITAT POLITÈCNICA  
DE CATALUNYA  
BARCELONATECH

In cooperation with  
CIMNE<sup>CS</sup>

## Advancing Fractional Calculus: A Fixed-Point Strategy for Nonlinear Caputo-Hadamard Equations

Muath Awadalla<sup>1</sup> and Manigandan Murugesan<sup>2,3,\*</sup>

<sup>1</sup>Department of Mathematics and Statistics, College of Science, King Faisal University, Hofuf, Al Ahsa, Saudi Arabia

<sup>2</sup>Center for Nonlinear and Complex Networks, SRM TRP Engineering College, Tiruchirapalli, Tamil Nadu, India

<sup>3</sup>Center for Research, Easwari Engineering College, Chennai, Tamil Nadu, India

### ABSTRACT

This paper investigates the existence, uniqueness, and Ulam–Hyers stability of solutions for a coupled system of nonlinear Caputo–Hadamard fractional differential equations in Banach spaces. By reformulating the boundary value problem into an equivalent integral system via the Hadamard fractional integral operator, sufficient conditions for existence and uniqueness are established using Krasnoselskii’s and Banach’s fixed-point theorems. Within the same functional framework, Ulam–Hyers stability results are derived for the proposed system. The theoretical analysis provides a consistent and unified approach for studying nonlinear coupled fractional systems with nonlocal operators, and the validity of the assumptions is illustrated through representative examples.

### OPEN ACCESS

**Received:** 14/01/2026

**Accepted:** 12/02/2026

**Published:** 16/04/2026

### DOI

10.23967/j.rimni.2026.10.79081

### Keywords:

Caputo–Hadamard fractional derivative  
Krasnoselskii fixed-point theorem  
Banach fixed-point theorem  
existence and uniqueness  
Ulam–Hyers stability  
fractional differential systems  
**MSC:** 34B15, 34B18, 34B37, 58E30.

## 1 Introduction

Fractional differential equations have become an essential tool for modeling complex phenomena with memory and hereditary effects, extending classical integer-order models in fields such as viscoelasticity, control theory, and biological systems [1–7]. Among various fractional operators, the Hadamard derivative, defined via a logarithmic kernel, is particularly suitable for problems on semi-infinite domains [8–10]. Its Caputo-type modification, the Caputo-Hadamard derivative, incorporates

\*Correspondence: Manigandan Murugesan (yogimani22@gmail.com). This is an article distributed under the terms of the Creative Commons BY-NC-SA license

initial conditions in a physically meaningful way, making it applicable to a wider range of practical problems [11–14].

Recent studies on boundary value problems involving Caputo-Hadamard derivatives have extended the analysis to various functional frameworks, including Banach and  $L^p$ -spaces, and have addressed numerical approximation, stability, and controllability aspects [15–20]. Fractional calculus has also found significant applications in optimization and machine learning, where fractional-order operators are used to design algorithms with enhanced convergence and robustness [21–24].

In [25], the authors studied the existence of solutions for a fractional boundary value problem:

$$\begin{cases} {}^c\mathcal{D}_1^{\gamma_1} \Phi(\zeta) = \omega_1(\zeta, \Phi(\zeta)), & \zeta \in [1, e], \\ \mathbb{A}_1\Phi(1) + \mathbb{A}_2\Phi'(1) = 0, & \mathbb{A}_3\Phi(e) + \mathbb{A}_4\Phi'(e) = \mathbb{A}_5, \end{cases} \quad (1)$$

where  ${}^c\mathcal{D}^{\gamma_1}$  denotes the Caputo fractional derivative.

Motivated by such works, this paper investigates a coupled system of nonlinear Caputo-Hadamard fractional differential equations:

$$\begin{cases} {}^{CH}\mathcal{D}_1^{\gamma_1} [\Phi(\zeta) - H\mathcal{J}_1^{\eta_1} \mathcal{H}_1(\zeta, \Phi(\zeta), \Omega(\zeta))] = \omega_1(\zeta, \Phi(\zeta), \Omega(\zeta)), & \zeta \in [1, e], \\ {}^{CH}\mathcal{D}_1^{\gamma_2} [\Omega(\zeta) - H\mathcal{J}_1^{\eta_2} \mathcal{H}_2(\zeta, \Phi(\zeta), \Omega(\zeta))] = \omega_2(\zeta, \Phi(\zeta), \Omega(\zeta)), & \zeta \in [1, e], \\ \mathbb{A}_1\Phi(1) + \mathbb{A}_2\Phi'(1) = 0, & \mathbb{A}_3\Phi(e) + \mathbb{A}_4\Phi'(e) = \mathbb{A}_5, \\ \mathbb{B}_1\Omega(1) + \mathbb{B}_2\Omega'(1) = 0, & \mathbb{B}_3\Omega(e) + \mathbb{B}_4\Omega'(e) = \mathbb{B}_5. \end{cases} \quad (2)$$

Here,  $1 < \gamma_1, \gamma_2 \leq 2$ , and  $\mathbb{A}_i, \mathbb{B}_j \in \mathbb{R}$  are real constants subject to standard non-degeneracy conditions, while  $\mathcal{H}_1, \mathcal{H}_2, \omega_1, \omega_2$  are given continuous functions.

The analysis proceeds by reformulating (2) into an equivalent integral system using the Hadamard fractional integral operator. Existence and uniqueness results are then established by combining Banach's contraction principle with Krasnoselskii's fixed-point theorem, the latter being applicable due to the compactness properties ensured by the Arzelà-Ascoli theorem. In the same functional setting, Ulam-Hyers stability criteria are derived for the coupled system. The consideration of distinct fractional derivative orders  $\gamma_i$  and integral orders  $\eta_i$  increases modeling flexibility but also introduces analytical challenges that are addressed under appropriate assumptions.

The paper is organized as follows. Section 2 collects necessary definitions, lemmas, and statements of the fixed-point theorems used. Section 3 presents the main existence and uniqueness theorems. Section 4 analyzes Ulam-Hyers stability. Section 5 provides illustrative examples, and Section 6 concludes the work.

## 2 Preliminaries

This section presents the definitions, theorems, and auxiliary results that form the foundation of our analysis. We begin with fractional calculus concepts, followed by the fixed-point theorems and stability definitions used throughout the paper.

### 2.1 Fractional Calculus Definitions

We recall the definitions of Hadamard fractional integral and Caputo-Hadamard fractional derivative, which are essential for our problem formulation.

**Definition 1 (Hadamard fractional integral [1]):** The Hadamard fractional integral of order  $\gamma > 0$  for a function  $h : (1, \infty) \rightarrow \mathbb{R}$  is defined as:

$$H\mathbb{J}_{1+}^{\gamma} h(\zeta) = \frac{1}{\Gamma(\gamma)} \int_1^{\zeta} \left( \log \frac{\zeta}{s} \right)^{\gamma-1} \frac{h(s)}{s} ds.$$

**Definition 2 (Caputo-Hadamard fractional derivative [8]):** Let  $\gamma > 0$ ,  $n = \lceil \gamma \rceil$ , and  $\delta = s \frac{d}{ds}$ . For  $\omega(\zeta) \in AC_{\delta}^n[a, b]$ , where  $0 < a < b < \infty$  and

$$AC_{\delta}^n[a, b] = \left\{ \omega : [a, b] \rightarrow \mathbb{R} \mid \delta^{n-1} \omega(\zeta) \in AC[a, b] \right\},$$

the Caputo-Hadamard fractional derivative of order  $\gamma$  is defined as:

$${}^{CH}D_{1+}^{\gamma} \omega(\zeta) = \frac{1}{\Gamma(n - \gamma)} \int_1^{\zeta} \left( \ln \frac{\zeta}{s} \right)^{n-\gamma-1} \delta^n \omega(s) \frac{ds}{s}.$$

**Lemma 1 (Solution of homogeneous equation [8]):** Let  $\gamma > 0$ ,  $n = \lceil \gamma \rceil$ . If  $\omega \in AC_{\delta}^n[a, b]$ , then the general solution of the equation

$${}^{CH}D_{1+}^{\gamma} \omega(\zeta) = 0$$

is given by

$$\omega(\zeta) = \sum_{i=0}^{n-1} c_i (\ln \zeta - \ln a)^i,$$

where  $c_i \in \mathbb{R}$  for  $i = 0, 1, \dots, n - 1$ . For the nonhomogeneous case, if  $z(\zeta) \in AC_{\delta}^n[a, b]$ , then

$$H\mathbb{J}_{1+}^{n-\gamma} ({}^{CH}D_{1+}^{\gamma} \omega) (\zeta) = z(\zeta) - \sum_{i=0}^{n-1} c_i (\ln \zeta - \ln a)^i.$$

## 2.2 Fixed-Point Theorems

The following fixed-point theorems are fundamental to our existence and uniqueness analysis.

**Theorem 1 (Banach Contraction Principle [2]):** Let  $(X, d)$  be a complete metric space and  $T : X \rightarrow X$  be a contraction mapping, i.e., there exists  $k \in [0, 1)$  such that

$$d(Tx, Ty) \leq k d(x, y) \quad \text{for all } x, y \in X.$$

Then  $T$  has a unique fixed point in  $X$ .

**Theorem 2 (Krasnoselskii's Fixed Point Theorem [26–28]):** Let  $M$  be a closed, bounded, convex, nonempty subset of a Banach space  $X$ . Suppose that  $A$  and  $B$  are operators mapping  $M$  into  $X$  such that:

1.  $Ax + Bx \in M$  for all  $x \in M$ ,
2.  $A$  is a contraction mapping,
3.  $B$  is compact and continuous.

Then there exists  $z \in M$  such that  $z = Az + Bz$ .

**Theorem 3 (Arzelà-Ascoli Theorem):** *A subset  $F$  of the space  $C([a, b], \mathbb{R})$  equipped with the supremum norm is relatively compact if and only if:*

1.  $F$  is uniformly bounded,
2.  $F$  is equicontinuous.

**Remark 1:** *The choice of the Arzelà-Ascoli theorem over other compactness criteria (such as Kolmogorov's criterion in measure spaces) is motivated by our working framework: the solution operators are naturally defined on the space of continuous functions  $C([1, e], \mathbb{R})$ , where pointwise boundedness and equicontinuity are the appropriate conditions for compactness.*

**Remark 2 (Analytical challenges):** *The main analytical difficulties in studying system (1.2) arise from three aspects: (i) the coupled structure involving both  $\Phi$  and  $\Omega$ , which requires handling product spaces; (ii) the presence of distinct fractional derivative orders  $\gamma_i$  and integral orders  $\eta_i$ , which complicates the estimation of norm bounds; and (iii) the combination of Hadamard integrals with Caputo-Hadamard derivatives, which demands careful manipulation of logarithmic kernels.*

### 2.3 Stability Definitions

The following definitions concern the Ulam-Hyers type stability of fractional differential systems.

**Definition 3 (Ulam-Hyers stability [13,14,29]):** *Let  $\epsilon = (\epsilon_1, \epsilon_2) > 0$ . The system (1.2) is said to be Ulam-Hyers stable if there exists a constant  $c = (c_1, c_2) > 0$  such that, for every solution  $(\Phi, \Omega) \in C([1, e], \mathbb{R}) \times C([1, e], \mathbb{R})$  of the inequalities*

$$\begin{aligned} |{}^{CH}D_{1+}^{\gamma_1}[\Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1}\mathcal{H}_1(\zeta, \Phi, \Omega)] - \omega_1(\zeta, \Phi, \Omega)| &\leq \epsilon_1, \\ |{}^{CH}D_{1+}^{\gamma_2}[\Omega(\zeta) - H\mathcal{J}_{1+}^{\eta_2}\mathcal{H}_2(\zeta, \Phi, \Omega)] - \omega_2(\zeta, \Phi, \Omega)| &\leq \epsilon_2, \end{aligned}$$

*there exists a unique solution  $(\hat{\Phi}, \hat{\Omega})$  of (1.2) satisfying*

$$\|(\Phi, \Omega) - (\hat{\Phi}, \hat{\Omega})\| \leq c\epsilon, \quad \zeta \in [1, e],$$

*where  $\|\cdot\|$  denotes an appropriate norm on the product space.*

**Definition 4 (Generalized Ulam-Hyers stability [29,30]):** *The system (1.2) is called generalized Ulam-Hyers stable if there exists a function  $\Psi \in C(\mathbb{R}^2, \mathbb{R}^2)$  with  $\Psi(0, 0) = (0, 0)$  such that for every solution  $(\Phi, \Omega)$  of the inequalities in Definition 3, there exists a solution  $(\hat{\Phi}, \hat{\Omega})$  of (1.2) with*

$$\|(\Phi, \Omega) - (\hat{\Phi}, \hat{\Omega})\| \leq \Psi(\epsilon_1, \epsilon_2).$$

**Definition 5 (Ulam-Hyers-Rassias stability [29]):** *Let  $\delta = (\delta_1, \delta_2) \in C([1, e], \mathbb{R}^2)$  be positive functions. The system (1.2) is Ulam-Hyers-Rassias stable with respect to  $\delta$  if there exists a constant  $c = (c_1, c_2) > 0$  such that, for every  $\epsilon = (\epsilon_1, \epsilon_2) > 0$  and every solution  $(\Phi, \Omega)$  of*

$$\begin{aligned} |{}^{CH}D_{1+}^{\gamma_1}[\Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1}\mathcal{H}_1(\zeta, \Phi, \Omega)] - \omega_1(\zeta, \Phi, \Omega)| &\leq \epsilon_1\delta_1(\zeta), \\ |{}^{CH}D_{1+}^{\gamma_2}[\Omega(\zeta) - H\mathcal{J}_{1+}^{\eta_2}\mathcal{H}_2(\zeta, \Phi, \Omega)] - \omega_2(\zeta, \Phi, \Omega)| &\leq \epsilon_2\delta_2(\zeta), \end{aligned}$$

*there exists a solution  $(\hat{\Phi}, \hat{\Omega})$  of (1.2) satisfying*

$$\|(\Phi, \Omega) - (\hat{\Phi}, \hat{\Omega})\| \leq c\delta(\zeta).$$

**Remark 3:** Unlike Lyapunov stability, which focuses on asymptotic behavior of trajectories, Ulam-Hyers stability deals with the approximation of solutions when the equation or its data are slightly perturbed. This qualitative stability concept is more suitable for our functional-analytic framework in Banach spaces.

### 3 Auxiliary Lemma: Integral Representation

**Lemma 2 (Integral representation for the linear system):** Let  $Q_1, Q_2, W_1, W_2 \in C([1, e], \mathbb{R})$ . Consider the linear coupled system

$$\begin{cases} {}^{CH}D_{1+}^{\gamma_1}[\Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1}Q_1(\zeta)] = Q_2(\zeta), & \zeta \in [1, e], \\ {}^{CH}D_{1+}^{\gamma_2}[\Omega(\zeta) - H\mathcal{J}_{1+}^{\eta_2}W_1(\zeta)] = W_2(\zeta), & \zeta \in [1, e], \\ \mathbb{A}_1\Phi(1) + \mathbb{A}_2\Phi'(1) = 0, & \mathbb{A}_3\Phi(e) + \mathbb{A}_4\Phi'(e) = \mathbb{A}_5, \\ \mathbb{B}_1\Omega(1) + \mathbb{B}_2\Omega'(1) = 0, & \mathbb{B}_3\Omega(e) + \mathbb{B}_4\Omega'(e) = \mathbb{B}_5, \end{cases} \quad (3)$$

where  $1 < \gamma_1, \gamma_2 \leq 2$  and  $\eta_1, \eta_2 > 1$ . Assume  $\mathbb{A}_1 \neq 0, \mathbb{B}_1 \neq 0$  and

$$\lambda_1 := \mathbb{A}_1\mathbb{A}_3e - \mathbb{A}_2\mathbb{A}_3e + \mathbb{A}_1\mathbb{A}_4 \neq 0, \quad \lambda_2 := \mathbb{B}_1\mathbb{B}_3e - \mathbb{B}_2\mathbb{B}_3e + \mathbb{B}_1\mathbb{B}_4 \neq 0.$$

(The restriction  $\eta_i > 1$  is imposed to ensure the differentiability required by the boundary conditions involving first derivatives.) Then the unique solution  $(\Phi, \Omega) \in C^1([1, e], \mathbb{R}) \times C^1([1, e], \mathbb{R})$  of (3) is given by

$$\begin{aligned} \Phi(\zeta) &= H\mathcal{J}_{1+}^{\gamma_1}Q_2(\zeta) + H\mathcal{J}_{1+}^{\eta_1}Q_1(\zeta) \\ &\quad - \frac{e(\mathbb{A}_2 - \mathbb{A}_1 \ln \zeta)}{\lambda_1} \left[ \mathbb{A}_5 - \mathbb{A}_3(H\mathcal{J}_{1+}^{\gamma_1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1}Q_1(e)) \right. \\ &\quad \left. - \frac{\mathbb{A}_4}{e}(H\mathcal{J}_{1+}^{\gamma_1-1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1-1}Q_1(e)) \right], \end{aligned} \quad (4)$$

and

$$\begin{aligned} \Omega(\zeta) &= H\mathcal{J}_{1+}^{\gamma_2}W_2(\zeta) + H\mathcal{J}_{1+}^{\eta_2}W_1(\zeta) \\ &\quad - \frac{e(\mathbb{B}_2 - \mathbb{B}_1 \ln \zeta)}{\lambda_2} \left[ \mathbb{B}_5 - \mathbb{B}_3(H\mathcal{J}_{1+}^{\gamma_2}W_2(e) + H\mathcal{J}_{1+}^{\eta_2}W_1(e)) \right. \\ &\quad \left. - \frac{\mathbb{B}_4}{e}(H\mathcal{J}_{1+}^{\gamma_2-1}W_2(e) + H\mathcal{J}_{1+}^{\eta_2-1}W_1(e)) \right], \end{aligned} \quad (5)$$

where the Hadamard fractional integral is defined as

$$H\mathcal{J}_{1+}^{\alpha}f(\zeta) = \frac{1}{\Gamma(\alpha)} \int_1^{\zeta} \left( \ln \frac{\zeta}{s} \right)^{\alpha-1} f(s) \frac{ds}{s}, \quad \alpha > 0.$$

**Proof.** We prove the representation for  $\Phi$ ; the derivation for  $\Omega$  follows identical steps with the replacements  $(\mathbb{A}_i, \lambda_1, \gamma_1, \eta_1, Q_1, Q_2) \mapsto (\mathbb{B}_i, \lambda_2, \gamma_2, \eta_2, W_1, W_2)$ .

From the first equation of (3),

$${}^{CH}D_{1+}^{\gamma_1}[\Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1}Q_1(\zeta)] = Q_2(\zeta), \quad \zeta \in [1, e].$$

Applying  $H\mathcal{J}_{1+}^{\gamma_1}$  to both sides and using the fundamental identity for  $1 < \gamma_1 \leq 2$ ,

$$H\mathcal{J}_{1+}^{\gamma_1}({}^{CH}D_{1+}^{\gamma_1}u)(\zeta) = u(\zeta) - c_0 - c_1 \ln \zeta,$$

we obtain

$$\Phi(\zeta) = H\mathcal{J}_{1+}^{\gamma_1}Q_2(\zeta) + H\mathcal{J}_{1+}^{\eta_1}Q_1(\zeta) + c_0 + c_1 \ln \zeta, \quad (6)$$

for constants  $c_0, c_1 \in \mathbb{R}$ .

Since  $\eta_1 > 1$  and  $1 < \gamma_1 \leq 2$ , the Hadamard integrals in (6) are continuously differentiable on  $[1, e]$ , hence  $\Phi \in C^1([1, e])$ . Differentiating and using the property

$$\frac{d}{d\zeta} H\mathcal{J}_{1+}^{\alpha}f(\zeta) = \frac{1}{\zeta} H\mathcal{J}_{1+}^{\alpha-1}f(\zeta) \quad (\alpha > 1),$$

we obtain

$$\Phi'(\zeta) = \frac{1}{\zeta} H\mathcal{J}_{1+}^{\gamma_1-1}Q_2(\zeta) + \frac{1}{\zeta} H\mathcal{J}_{1+}^{\eta_1-1}Q_1(\zeta) + \frac{c_1}{\zeta}. \quad (7)$$

Evaluating (6) and (7) at  $\zeta = 1$  gives  $\Phi(1) = c_0$  and  $\Phi'(1) = c_1$ . Therefore,

$$\mathbb{A}_1\Phi(1) + \mathbb{A}_2\Phi'(1) = 0 \implies \mathbb{A}_1c_0 + \mathbb{A}_2c_1 = 0.$$

With  $\mathbb{A}_1 \neq 0$ , this yields

$$c_0 = -\frac{\mathbb{A}_2}{\mathbb{A}_1}c_1. \quad (8)$$

From (6) and (7) at  $\zeta = e$ :

$$\Phi(e) = H\mathcal{J}_{1+}^{\gamma_1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1}Q_1(e) + c_0 + c_1,$$

$$\Phi'(e) = \frac{1}{e} \left( H\mathcal{J}_{1+}^{\gamma_1-1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1-1}Q_1(e) \right) + \frac{c_1}{e}.$$

Substituting these into  $\mathbb{A}_3\Phi(e) + \mathbb{A}_4\Phi'(e) = \mathbb{A}_5$  and using (8):

$$\begin{aligned} \mathbb{A}_5 &= \mathbb{A}_3 \left( H\mathcal{J}_{1+}^{\gamma_1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1}Q_1(e) + c_1 - \frac{\mathbb{A}_2}{\mathbb{A}_1}c_1 \right) \\ &\quad + \mathbb{A}_4 \left( \frac{1}{e} \left( H\mathcal{J}_{1+}^{\gamma_1-1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1-1}Q_1(e) \right) + \frac{c_1}{e} \right) \\ &= \mathbb{A}_3 \left( H\mathcal{J}_{1+}^{\gamma_1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1}Q_1(e) \right) \\ &\quad + \frac{\mathbb{A}_4}{e} \left( H\mathcal{J}_{1+}^{\gamma_1-1}Q_2(e) + H\mathcal{J}_{1+}^{\eta_1-1}Q_1(e) \right) \\ &\quad + c_1 \left( \mathbb{A}_3 \left( 1 - \frac{\mathbb{A}_2}{\mathbb{A}_1} \right) + \frac{\mathbb{A}_4}{e} \right). \end{aligned}$$

The coefficient of  $c_1$  simplifies as

$$\mathbb{A}_3 \left( 1 - \frac{\mathbb{A}_2}{\mathbb{A}_1} \right) + \frac{\mathbb{A}_4}{e} = \frac{1}{\mathbb{A}_1 e} \left( \mathbb{A}_1 \mathbb{A}_3 e - \mathbb{A}_2 \mathbb{A}_3 e + \mathbb{A}_1 \mathbb{A}_4 \right) = \frac{\lambda_1}{\mathbb{A}_1 e}.$$

Since  $\lambda_1 \neq 0$ ,

$$c_1 = \frac{\mathbb{A}_1 e}{\lambda_1} \left[ \mathbb{A}_5 - \mathbb{A}_3 \left( H\mathcal{J}_{1+}^{\gamma_1} \mathcal{Q}_2(e) + H\mathcal{J}_{1+}^{\eta_1} \mathcal{Q}_1(e) \right) - \frac{\mathbb{A}_4}{e} \left( H\mathcal{J}_{1+}^{\gamma_1-1} \mathcal{Q}_2(e) + H\mathcal{J}_{1+}^{\eta_1-1} \mathcal{Q}_1(e) \right) \right].$$

From (8),

$$c_0 = -\frac{\mathbb{A}_2}{\mathbb{A}_1} c_1 = -\frac{\mathbb{A}_2 e}{\lambda_1} \left[ \mathbb{A}_5 - \mathbb{A}_3 \left( H\mathcal{J}_{1+}^{\gamma_1} \mathcal{Q}_2(e) + H\mathcal{J}_{1+}^{\eta_1} \mathcal{Q}_1(e) \right) - \frac{\mathbb{A}_4}{e} \left( H\mathcal{J}_{1+}^{\gamma_1-1} \mathcal{Q}_2(e) + H\mathcal{J}_{1+}^{\eta_1-1} \mathcal{Q}_1(e) \right) \right].$$

Now substitute  $c_0$  and  $c_1$  into (6). Observing that

$$c_0 + c_1 \ln \zeta = c_1 \left( \ln \zeta - \frac{\mathbb{A}_2}{\mathbb{A}_1} \right) = -\frac{\mathbb{A}_2 - \mathbb{A}_1 \ln \zeta}{\mathbb{A}_1} c_1,$$

and using the expression for  $c_1$ , we obtain precisely (4).

To verify that (4) satisfies the first equation of (3), note from (6) that

$$\Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1} \mathcal{Q}_1(\zeta) = H\mathcal{J}_{1+}^{\gamma_1} \mathcal{Q}_2(\zeta) + c_0 + c_1 \ln \zeta.$$

Applying  ${}^{CH}D_{1+}^{\gamma_1}$  to both sides and using  ${}^{CH}D_{1+}^{\gamma_1} H\mathcal{J}_{1+}^{\gamma_1} f = f$  together with  ${}^{CH}D_{1+}^{\gamma_1} (c_0 + c_1 \ln \zeta) = 0$  for  $1 < \gamma_1 \leq 2$ , we recover

$${}^{CH}D_{1+}^{\gamma_1} [\Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1} \mathcal{Q}_1(\zeta)] = \mathcal{Q}_2(\zeta).$$

The boundary conditions are satisfied by construction because  $c_0$  and  $c_1$  were uniquely determined from them. Hence, (4) is the unique solution component  $\Phi$ . The same reasoning applied to the second equation yields (5) for  $\Omega$ .  $\square$

**Remark 4:** The explicit Formulas (4) and (5) reveal the structure of the solution operator: a local part consisting of Hadamard integrals from 1 to  $\zeta$ , and a non-local part that incorporates the boundary data via integrals over the whole interval  $[1, e]$ . The non-degeneracy conditions  $\lambda_1 \neq 0$  and  $\lambda_2 \neq 0$  guarantee that the constants  $c_0, c_1$  (and their analogues for  $\Omega$ ) are uniquely determined, which makes the linear boundary-value problem well-posed. This representation will be used in the next section to reformulate the original nonlinear problem as a fixed-point equation.

## 4 Main Results

### 4.1 Functional Setting and Operator Formulation

Let us introduce the Banach space

$$\mathcal{X} = C([1, e], \mathbb{R}) \times C([1, e], \mathbb{R})$$

equipped with the norm

$$\|(\Phi, \Omega)\| = \|\Phi\| + \|\Omega\|, \quad \|\Phi\| = \max_{\zeta \in [1, e]} |\Phi(\zeta)|, \quad \|\Omega\| = \max_{\zeta \in [1, e]} |\Omega(\zeta)|.$$

Using the integral representation given in Lemma 2, we define an operator  $\mathcal{T} : \mathcal{X} \rightarrow \mathcal{X}$  by  $\mathcal{T}(\Phi, \Omega) = (\mathcal{T}_1(\Phi, \Omega), \mathcal{T}_2(\Phi, \Omega))$ , where

$$\begin{aligned} \mathcal{T}_1(\Phi, \Omega)(\zeta) &= H\mathcal{J}_{1+}^{\gamma_1} \omega_1(\zeta, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_1} \mathcal{H}_1(\zeta, \Phi, \Omega) \\ &\quad - \frac{e(\mathbb{A}_2 - \mathbb{A}_1 \ln \zeta)}{\lambda_1} \left[ \mathbb{A}_5 - \mathbb{A}_3 (H\mathcal{J}_{1+}^{\gamma_1} \omega_1(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_1} \mathcal{H}_1(e, \Phi, \Omega)) \right. \\ &\quad \left. - \frac{\mathbb{A}_4}{e} (H\mathcal{J}_{1+}^{\gamma_1-1} \omega_1(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_1-1} \mathcal{H}_1(e, \Phi, \Omega)) \right], \end{aligned} \quad (9)$$

and

$$\begin{aligned} \mathcal{T}_2(\Phi, \Omega)(\zeta) &= H\mathcal{J}_{1+}^{\gamma_2} \omega_2(\zeta, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_2} \mathcal{H}_2(\zeta, \Phi, \Omega) \\ &\quad - \frac{e(\mathbb{B}_2 - \mathbb{B}_1 \ln \zeta)}{\lambda_2} \left[ \mathbb{B}_5 - \mathbb{B}_3 (H\mathcal{J}_{1+}^{\gamma_2} \omega_2(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_2} \mathcal{H}_2(e, \Phi, \Omega)) \right. \\ &\quad \left. - \frac{\mathbb{B}_4}{e} (H\mathcal{J}_{1+}^{\gamma_2-1} \omega_2(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_2-1} \mathcal{H}_2(e, \Phi, \Omega)) \right]. \end{aligned} \quad (10)$$

Here, the Hadamard fractional integrals are understood pointwise, i.e.,

$$H\mathcal{J}_{1+}^{\alpha} f(\zeta, \Phi, \Omega) = \frac{1}{\Gamma(\alpha)} \int_1^{\zeta} \left( \ln \frac{\zeta}{s} \right)^{\alpha-1} f(s, \Phi(s), \Omega(s)) \frac{ds}{s}, \quad \alpha > 0.$$

A fixed point of  $\mathcal{T}$  corresponds precisely to a solution of the original nonlinear system (1). Consequently, establishing existence and uniqueness of solutions for (1) is equivalent to proving that  $\mathcal{T}$  possesses a unique fixed point in  $\mathcal{X}$ .

#### 4.2 Key Constants and Assumptions

We shall work under the following hypotheses:

(A<sub>1</sub>) The functions  $\mathcal{H}_1, \mathcal{H}_2, \omega_1, \omega_2 : [1, e] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous and satisfy Lipschitz conditions with respect to the second and third variables: there exist constants  $L_{\mathcal{H}_1}, L_{\mathcal{H}_2}, L_{\omega_1}, L_{\omega_2} > 0$  such that for all  $\zeta \in [1, e]$  and  $(\Phi_1, \Omega_1), (\Phi_2, \Omega_2) \in \mathbb{R}^2$ ,

$$|\mathcal{H}_i(\zeta, \Phi_1, \Omega_1) - \mathcal{H}_i(\zeta, \Phi_2, \Omega_2)| \leq L_{\mathcal{H}_i} (|\Phi_1 - \Phi_2| + |\Omega_1 - \Omega_2|),$$

$$|\omega_i(\zeta, \Phi_1, \Omega_1) - \omega_i(\zeta, \Phi_2, \Omega_2)| \leq L_{\omega_i} (|\Phi_1 - \Phi_2| + |\Omega_1 - \Omega_2|), \quad i = 1, 2.$$

(A<sub>2</sub>) The following boundedness conditions hold:

$$\mathcal{M}_1 := \max_{\zeta \in [1, e]} |\omega_1(\zeta, 0, 0)| < \infty, \quad \mathcal{M}_2 := \max_{\zeta \in [1, e]} |\mathcal{H}_1(\zeta, 0, 0)| < \infty,$$

$$\mathcal{N}_1 := \max_{\zeta \in [1, e]} |\omega_2(\zeta, 0, 0)| < \infty, \quad \mathcal{N}_2 := \max_{\zeta \in [1, e]} |\mathcal{H}_2(\zeta, 0, 0)| < \infty.$$

(A<sub>3</sub>) There exist continuous functions  $\psi_1, \psi_2, \psi_3, \psi_4 \in C([1, e], \mathbb{R}^+)$  such that

$$|\omega_1(\zeta, \Phi, \Omega)| \leq \psi_1(\zeta),$$

$$|\mathcal{H}_1(\zeta, \Phi, \Omega)| \leq \psi_2(\zeta),$$

$$|\omega_2(\zeta, \Phi, \Omega)| \leq \psi_3(\zeta),$$

$$|\mathcal{H}_2(\zeta, \Phi, \Omega)| \leq \psi_4(\zeta),$$

for all  $\zeta \in [1, e]$  and  $(\Phi, \Omega) \in \mathbb{R}^2$ .

For convenience we introduce the aggregated Lipschitz constants

$$\mathcal{L}_1 = L_{\mathcal{H}_1} + L_{\omega_1}, \quad \mathcal{L}_2 = L_{\mathcal{H}_2} + L_{\omega_2},$$

and the quantities

$$\mathcal{K}_1 = \frac{1}{\Gamma(\gamma_1 + 1)} + \frac{1}{\Gamma(\eta_1 + 1)} + \frac{|\mathbb{A}_2 \mathbb{A}_3| e}{|\lambda_1|} \left( \frac{1}{\Gamma(\gamma_1 + 1)} + \frac{1}{\Gamma(\eta_1 + 1)} \right) + \frac{|\mathbb{A}_2 \mathbb{A}_4|}{|\lambda_1|} \left( \frac{1}{\Gamma(\gamma_1)} + \frac{1}{\Gamma(\eta_1)} \right), \quad (11)$$

$$\mathcal{K}_2 = \frac{1}{\Gamma(\gamma_2 + 1)} + \frac{1}{\Gamma(\eta_2 + 1)} + \frac{|\mathbb{B}_2 \mathbb{B}_3| e}{|\lambda_2|} \left( \frac{1}{\Gamma(\gamma_2 + 1)} + \frac{1}{\Gamma(\eta_2 + 1)} \right) + \frac{|\mathbb{B}_2 \mathbb{B}_4|}{|\lambda_2|} \left( \frac{1}{\Gamma(\gamma_2)} + \frac{1}{\Gamma(\eta_2)} \right). \quad (12)$$

These constants will play a crucial role in the subsequent contraction and boundedness estimates.

#### 4.3 Theorem 4: Existence and Uniqueness via Banach's Fixed Point Theorem

**Theorem 4 (Existence and uniqueness):** Assume that conditions  $(A_1)$  and  $(A_2)$  are satisfied and that

$$\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2 < 1. \quad (13)$$

Then the system (1) possesses a unique solution in  $\mathcal{X}$ .

**Proof:** We shall prove that the operator  $\mathcal{T}$  defined by (9)–(10) is a contraction on a suitable closed ball of  $\mathcal{X}$ . Once this is established, Banach's contraction principle (Theorem 1) guarantees the existence of a unique fixed point, which is precisely the desired solution.

**1. Self-mapping on a closed ball.** Choose  $\varepsilon > 0$  sufficiently large so that

$$\varepsilon \geq [\mathcal{L}_1 \varepsilon + (\mathcal{M}_1 + \mathcal{M}_2)] \mathcal{K}_1 + [\mathcal{L}_2 \varepsilon + (\mathcal{N}_1 + \mathcal{N}_2)] \mathcal{K}_2 + e \left( \frac{|\mathbb{A}_2 \mathbb{A}_5|}{|\lambda_1|} + \frac{|\mathbb{B}_2 \mathbb{B}_5|}{|\lambda_2|} \right). \quad (14)$$

Such an  $\varepsilon$  exists because the right-hand side of (14) is an affine function of  $\varepsilon$  with slope  $\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2 < 1$  by (13). Define the closed ball

$$B_\varepsilon = \{(\Phi, \Omega) \in \mathcal{X} : \|(\Phi, \Omega)\| \leq \varepsilon\}.$$

Take  $(\Phi, \Omega) \in B_\varepsilon$ . For any  $s \in [1, e]$ , using  $(A_1)$  and  $(A_2)$  we have

$$\begin{aligned} |\omega_1(s, \Phi, \Omega)| &\leq |\omega_1(s, \Phi, \Omega) - \omega_1(s, 0, 0)| + |\omega_1(s, 0, 0)| \\ &\leq L_{\omega_1} (|\Phi(s)| + |\Omega(s)|) + \mathcal{M}_1 \\ &\leq L_{\omega_1} \varepsilon + \mathcal{M}_1, \end{aligned}$$

and similarly

$$|\mathcal{H}_1(s, \Phi, \Omega)| \leq L_{\mathcal{H}_1} \varepsilon + \mathcal{M}_2.$$

Now estimate each term in (9). For the local integrals, using the elementary identity

$$\int_1^\zeta \left( \ln \frac{\zeta}{s} \right)^{\alpha-1} \frac{ds}{s} = \frac{(\ln \zeta)^\alpha}{\alpha}, \quad \alpha > 0,$$

we obtain

$$|H\mathcal{J}_{1+}^{\gamma_1}\omega_1(\zeta, \Phi, \Omega)| \leq \frac{(L_{\omega_1}\varepsilon + \mathcal{M}_1)}{\Gamma(\gamma_1 + 1)}(\ln \zeta)^{\gamma_1} \leq \frac{L_{\omega_1}\varepsilon + \mathcal{M}_1}{\Gamma(\gamma_1 + 1)},$$

since  $\ln \zeta \leq 1$  for  $\zeta \in [1, e]$ . Analogously,

$$|H\mathcal{J}_{1+}^{\eta_1}\mathcal{H}_1(\zeta, \Phi, \Omega)| \leq \frac{L_{\mathcal{H}_1}\varepsilon + \mathcal{M}_2}{\Gamma(\eta_1 + 1)}.$$

For the non-local part, observe first that  $|\mathbb{A}_2 - \mathbb{A}_1 \ln \zeta| \leq |\mathbb{A}_2| + |\mathbb{A}_1|$  for  $\zeta \in [1, e]$ . Moreover,

$$|H\mathcal{J}_{1+}^{\gamma_1}\omega_1(e, \Phi, \Omega)| \leq \frac{L_{\omega_1}\varepsilon + \mathcal{M}_1}{\Gamma(\gamma_1 + 1)}, \quad |H\mathcal{J}_{1+}^{\eta_1}\mathcal{H}_1(e, \Phi, \Omega)| \leq \frac{L_{\mathcal{H}_1}\varepsilon + \mathcal{M}_2}{\Gamma(\eta_1 + 1)},$$

and for the integrals of order  $\gamma_1 - 1$  and  $\eta_1 - 1$ ,

$$|H\mathcal{J}_{1+}^{\gamma_1-1}\omega_1(e, \Phi, \Omega)| \leq \frac{L_{\omega_1}\varepsilon + \mathcal{M}_1}{\Gamma(\gamma_1)}, \quad |H\mathcal{J}_{1+}^{\eta_1-1}\mathcal{H}_1(e, \Phi, \Omega)| \leq \frac{L_{\mathcal{H}_1}\varepsilon + \mathcal{M}_2}{\Gamma(\eta_1)}.$$

Assembling these estimates, we find

$$\begin{aligned} |\mathcal{T}_1(\Phi, \Omega)(\zeta)| &\leq \frac{L_{\omega_1}\varepsilon + \mathcal{M}_1}{\Gamma(\gamma_1 + 1)} + \frac{L_{\mathcal{H}_1}\varepsilon + \mathcal{M}_2}{\Gamma(\eta_1 + 1)} \\ &\quad + \frac{e(|\mathbb{A}_2| + |\mathbb{A}_1|)}{|\lambda_1|} \left[ |\mathbb{A}_5| + |\mathbb{A}_3| \left( \frac{L_{\omega_1}\varepsilon + \mathcal{M}_1}{\Gamma(\gamma_1 + 1)} + \frac{L_{\mathcal{H}_1}\varepsilon + \mathcal{M}_2}{\Gamma(\eta_1 + 1)} \right) \right. \\ &\quad \left. + \frac{|\mathbb{A}_4|}{e} \left( \frac{L_{\omega_1}\varepsilon + \mathcal{M}_1}{\Gamma(\gamma_1)} + \frac{L_{\mathcal{H}_1}\varepsilon + \mathcal{M}_2}{\Gamma(\eta_1)} \right) \right]. \end{aligned}$$

After straightforward algebraic manipulation this yields

$$|\mathcal{T}_1(\Phi, \Omega)(\zeta)| \leq [\mathcal{L}_1\varepsilon + (\mathcal{M}_1 + \mathcal{M}_2)]\mathcal{K}_1 + \frac{e|\mathbb{A}_2\mathbb{A}_5|}{|\lambda_1|}.$$

A completely analogous calculation for  $\mathcal{T}_2$  gives

$$|\mathcal{T}_2(\Phi, \Omega)(\zeta)| \leq [\mathcal{L}_2\varepsilon + (\mathcal{N}_1 + \mathcal{N}_2)]\mathcal{K}_2 + \frac{e|\mathbb{B}_2\mathbb{B}_5|}{|\lambda_2|}.$$

Consequently,

$$\begin{aligned} \|\mathcal{T}(\Phi, \Omega)\| &= \|\mathcal{T}_1(\Phi, \Omega)\| + \|\mathcal{T}_2(\Phi, \Omega)\| \\ &\leq [\mathcal{L}_1\varepsilon + (\mathcal{M}_1 + \mathcal{M}_2)]\mathcal{K}_1 + [\mathcal{L}_2\varepsilon + (\mathcal{N}_1 + \mathcal{N}_2)]\mathcal{K}_2 + e \left( \frac{|\mathbb{A}_2\mathbb{A}_5|}{|\lambda_1|} + \frac{|\mathbb{B}_2\mathbb{B}_5|}{|\lambda_2|} \right) \\ &\leq \varepsilon, \end{aligned}$$

where the last inequality is exactly condition (14). Hence  $\mathcal{T}(B_\varepsilon) \subset B_\varepsilon$ .

**2. Contraction property.** Take  $(\Phi_1, \Omega_1), (\Phi_2, \Omega_2) \in B_\varepsilon$ . Using the Lipschitz conditions  $(A_1)$ ,

$$\begin{aligned} |\omega_1(s, \Phi_1, \Omega_1) - \omega_1(s, \Phi_2, \Omega_2)| &\leq L_{\omega_1} (|\Phi_1(s) - \Phi_2(s)| + |\Omega_1(s) - \Omega_2(s)|), \\ |\mathcal{H}_1(s, \Phi_1, \Omega_1) - \mathcal{H}_1(s, \Phi_2, \Omega_2)| &\leq L_{\mathcal{H}_1} (|\Phi_1(s) - \Phi_2(s)| + |\Omega_1(s) - \Omega_2(s)|). \end{aligned}$$

Insert these estimates into (9). Proceeding as in part 1 and using the fact that

$$\max_{s \in [1, e]} (|\Phi_1(s) - \Phi_2(s)| + |\Omega_1(s) - \Omega_2(s)|) \leq \|\Phi_1 - \Phi_2\| + \|\Omega_1 - \Omega_2\|,$$

we obtain after evaluating all integrals

$$|\mathcal{T}_1(\Phi_1, \Omega_1)(\zeta) - \mathcal{T}_1(\Phi_2, \Omega_2)(\zeta)| \leq \mathcal{L}_1 \mathcal{K}_1 (\|\Phi_1 - \Phi_2\| + \|\Omega_1 - \Omega_2\|).$$

A parallel computation for the second component yields

$$|\mathcal{T}_2(\Phi_1, \Omega_1)(\zeta) - \mathcal{T}_2(\Phi_2, \Omega_2)(\zeta)| \leq \mathcal{L}_2 \mathcal{K}_2 (\|\Phi_1 - \Phi_2\| + \|\Omega_1 - \Omega_2\|).$$

Therefore,

$$\begin{aligned} \|\mathcal{T}(\Phi_1, \Omega_1) - \mathcal{T}(\Phi_2, \Omega_2)\| &\leq (\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2) (\|\Phi_1 - \Phi_2\| + \|\Omega_1 - \Omega_2\|) \\ &= (\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2) \|(\Phi_1, \Omega_1) - (\Phi_2, \Omega_2)\|. \end{aligned}$$

Condition (13) guarantees that  $\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2 < 1$ ; thus  $\mathcal{T}$  is a contraction on  $B_\varepsilon$ .

**3. Application of Banach's theorem and global uniqueness.** The closed ball  $B_\varepsilon$  is a complete metric space (as a closed subset of the complete space  $\mathcal{X}$ ). The operator  $\mathcal{T}$  maps  $B_\varepsilon$  into itself and is a contraction. By Banach's fixed point theorem (Theorem 1),  $\mathcal{T}$  possesses a unique fixed point in  $B_\varepsilon$ . This fixed point is a solution of the original system (1).

To see that uniqueness holds in the whole space  $\mathcal{X}$ , note that any solution  $(\Phi, \Omega) \in \mathcal{X}$  is necessarily bounded on  $[1, e]$  because the functions  $\mathcal{H}_i, \omega_i$  are continuous and the fractional integrals preserve boundedness. Hence every solution belongs to some ball  $B_\varepsilon$  with a sufficiently large radius  $\varepsilon$ . Since  $\mathcal{T}$  has exactly one fixed point in each such ball, the solution is unique globally in  $\mathcal{X}$ .  $\square$

#### 4.4 Theorem 5: Existence via Krasnoselskii's Fixed Point Theorem

**Theorem 5 (Existence):** Assume that conditions  $(A_1)$  and  $(A_3)$  are satisfied. If, in addition,

$$\mathcal{L}_1 \left( \frac{1}{\Gamma(\gamma_1 + 1)} + \frac{1}{\Gamma(\eta_1 + 1)} \right) < 1, \tag{15}$$

and

$$\mathcal{L}_2 \left( \frac{1}{\Gamma(\gamma_2 + 1)} + \frac{1}{\Gamma(\eta_2 + 1)} \right) < 1, \tag{16}$$

then the system (1) possesses at least one solution in  $\mathcal{X}$ .

**Proof.** We apply Krasnoselskii's fixed point theorem (Theorem 2) on a suitable closed, bounded, convex subset of  $\mathcal{X}$  by decomposing the operator  $\mathcal{T}$  as  $\mathcal{T} = \mathcal{A} + \mathcal{B}$ .

**Step 1: Decomposition.** Define  $\mathcal{A} = (\mathcal{A}_1, \mathcal{A}_2)$  and  $\mathcal{B} = (\mathcal{B}_1, \mathcal{B}_2)$  on  $\mathcal{X}$  by

$$\mathcal{A}_1(\Phi, \Omega)(\zeta) = H \mathcal{J}_{1+}^{\gamma_1} \omega_1(\zeta, \Phi, \Omega) + H \mathcal{J}_{1+}^{\eta_1} \mathcal{H}_1(\zeta, \Phi, \Omega),$$

$$\mathcal{A}_2(\Phi, \Omega)(\zeta) = H \mathcal{J}_{1+}^{\gamma_2} \omega_2(\zeta, \Phi, \Omega) + H \mathcal{J}_{1+}^{\eta_2} \mathcal{H}_2(\zeta, \Phi, \Omega),$$

and

$$\begin{aligned} \mathcal{B}_1(\Phi, \Omega)(\zeta) &= -\frac{e(\mathbb{A}_2 - \mathbb{A}_1 \ln \zeta)}{\lambda_1} \left[ \mathbb{A}_5 - \mathbb{A}_3 (H\mathcal{J}_{1+}^{\gamma_1} \omega_1(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_1} \mathcal{H}_1(e, \Phi, \Omega)) \right. \\ &\quad \left. - \frac{\mathbb{A}_4}{e} (H\mathcal{J}_{1+}^{\gamma_1-1} \omega_1(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_1-1} \mathcal{H}_1(e, \Phi, \Omega)) \right], \\ \mathcal{B}_2(\Phi, \Omega)(\zeta) &= -\frac{e(\mathbb{B}_2 - \mathbb{B}_1 \ln \zeta)}{\lambda_2} \left[ \mathbb{B}_5 - \mathbb{B}_3 (H\mathcal{J}_{1+}^{\gamma_2} \omega_2(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_2} \mathcal{H}_2(e, \Phi, \Omega)) \right. \\ &\quad \left. - \frac{\mathbb{B}_4}{e} (H\mathcal{J}_{1+}^{\gamma_2-1} \omega_2(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_2-1} \mathcal{H}_2(e, \Phi, \Omega)) \right]. \end{aligned}$$

Clearly,  $\mathcal{T} = \mathcal{A} + \mathcal{B}$ .

**Step 2: Choice of the invariant closed ball.** Let

$$r \geq 2 \max \left\{ (\|\psi_1\|_\infty + \|\psi_2\|_\infty) \mathcal{K}_1 + \frac{e|\mathbb{A}_2 \mathbb{A}_5|}{|\lambda_1|}, (\|\psi_3\|_\infty + \|\psi_4\|_\infty) \mathcal{K}_2 + \frac{e|\mathbb{B}_2 \mathbb{B}_5|}{|\lambda_2|} \right\},$$

where  $\|\psi_i\|_\infty = \max_{\zeta \in [1, e]} \psi_i(\zeta)$ . Consider the closed ball

$$\bar{D}_r = \{(\Phi, \Omega) \in \mathcal{X} : \|(\Phi, \Omega)\| \leq r\}.$$

For any  $(\Phi, \Omega) \in \bar{D}_r$ , assumption  $(A_3)$  yields the uniform bounds

$$\begin{aligned} |\omega_1(\zeta, \Phi, \Omega)| &\leq \|\psi_1\|_\infty, & |\mathcal{H}_1(\zeta, \Phi, \Omega)| &\leq \|\psi_2\|_\infty, \\ |\omega_2(\zeta, \Phi, \Omega)| &\leq \|\psi_3\|_\infty, & |\mathcal{H}_2(\zeta, \Phi, \Omega)| &\leq \|\psi_4\|_\infty. \end{aligned}$$

Using the same Hadamard-kernel estimates employed in the definition of  $\mathcal{K}_1, \mathcal{K}_2$  (with Lipschitz bounds replaced by the above uniform bounds), we get

$$\begin{aligned} \|\mathcal{T}_1(\Phi, \Omega)\| &\leq (\|\psi_1\|_\infty + \|\psi_2\|_\infty) \mathcal{K}_1 + \frac{e|\mathbb{A}_2 \mathbb{A}_5|}{|\lambda_1|}, \\ \|\mathcal{T}_2(\Phi, \Omega)\| &\leq (\|\psi_3\|_\infty + \|\psi_4\|_\infty) \mathcal{K}_2 + \frac{e|\mathbb{B}_2 \mathbb{B}_5|}{|\lambda_2|}. \end{aligned}$$

Consequently,

$$\|\mathcal{T}(\Phi, \Omega)\| = \|\mathcal{T}_1(\Phi, \Omega)\| + \|\mathcal{T}_2(\Phi, \Omega)\| \leq \frac{r}{2} + \frac{r}{2} = r,$$

which implies  $\mathcal{T}(\bar{D}_r) \subset \bar{D}_r$ . In particular,  $\mathcal{A}(\bar{D}_r) + \mathcal{B}(\bar{D}_r) \subset \bar{D}_r$ .

**Step 3:  $\mathcal{A}$  is a contraction on  $\bar{D}_r$ .** Let  $(\Phi_1, \Omega_1), (\Phi_2, \Omega_2) \in \bar{D}_r$ . By  $(A_1)$ ,

$$\begin{aligned} |\mathcal{A}_1(\Phi_1, \Omega_1)(\zeta) - \mathcal{A}_1(\Phi_2, \Omega_2)(\zeta)| &\leq \frac{L_{\omega_1}}{\Gamma(\gamma_1)} \int_1^\zeta \left( \ln \frac{\zeta}{s} \right)^{\gamma_1-1} (|\Phi_1(s) - \Phi_2(s)| + |\Omega_1(s) - \Omega_2(s)|) \frac{ds}{s} \\ &\quad + \frac{L_{\mathcal{H}_1}}{\Gamma(\eta_1)} \int_1^\zeta \left( \ln \frac{\zeta}{s} \right)^{\eta_1-1} (|\Phi_1(s) - \Phi_2(s)| + |\Omega_1(s) - \Omega_2(s)|) \frac{ds}{s}. \end{aligned}$$

Since  $\ln \zeta \leq 1$  on  $[1, e]$  and  $\Gamma(\alpha + 1) = \alpha \Gamma(\alpha)$ , we have for  $\alpha > 0$ ,

$$\frac{1}{\Gamma(\alpha)} \int_1^\zeta \left( \ln \frac{\zeta}{s} \right)^{\alpha-1} \frac{ds}{s} = \frac{(\ln \zeta)^\alpha}{\Gamma(\alpha + 1)} \leq \frac{1}{\Gamma(\alpha + 1)}.$$

Therefore,

$$|\mathcal{A}_1(\Phi_1, \Omega_1)(\zeta) - \mathcal{A}_1(\Phi_2, \Omega_2)(\zeta)| \leq \mathcal{L}_1 \left( \frac{1}{\Gamma(\gamma_1 + 1)} + \frac{1}{\Gamma(\eta_1 + 1)} \right) \|(\Phi_1, \Omega_1) - (\Phi_2, \Omega_2)\|.$$

Taking the supremum over  $\zeta \in [1, e]$  gives

$$\|\mathcal{A}_1(\Phi_1, \Omega_1) - \mathcal{A}_1(\Phi_2, \Omega_2)\| \leq \mathcal{L}_1 \left( \frac{1}{\Gamma(\gamma_1 + 1)} + \frac{1}{\Gamma(\eta_1 + 1)} \right) \|(\Phi_1, \Omega_1) - (\Phi_2, \Omega_2)\|.$$

Similarly,

$$\|\mathcal{A}_2(\Phi_1, \Omega_1) - \mathcal{A}_2(\Phi_2, \Omega_2)\| \leq \mathcal{L}_2 \left( \frac{1}{\Gamma(\gamma_2 + 1)} + \frac{1}{\Gamma(\eta_2 + 1)} \right) \|(\Phi_1, \Omega_1) - (\Phi_2, \Omega_2)\|.$$

Hence, letting

$$\rho = \max \left\{ \mathcal{L}_1 \left( \frac{1}{\Gamma(\gamma_1 + 1)} + \frac{1}{\Gamma(\eta_1 + 1)} \right), \mathcal{L}_2 \left( \frac{1}{\Gamma(\gamma_2 + 1)} + \frac{1}{\Gamma(\eta_2 + 1)} \right) \right\},$$

we obtain

$$\|\mathcal{A}(\Phi_1, \Omega_1) - \mathcal{A}(\Phi_2, \Omega_2)\| \leq \rho \|(\Phi_1, \Omega_1) - (\Phi_2, \Omega_2)\|.$$

By (15) and (16),  $\rho < 1$ , so  $\mathcal{A}$  is a contraction on  $\overline{D}_r$ .

**Step 4:  $\mathcal{B}$  is compact and continuous on  $\overline{D}_r$ .** We show the claim for  $\mathcal{B}_1$ ; the proof for  $\mathcal{B}_2$  is analogous. Write

$$\mathcal{B}_1(\Phi, \Omega)(\zeta) = \theta(\zeta) \Psi(\Phi, \Omega),$$

where  $\theta(\zeta) = -\frac{e(\mathbb{A}_2 - \mathbb{A}_1 \ln \zeta)}{\lambda_1}$  is a fixed  $C^1$  function on  $[1, e]$ , and

$$\begin{aligned} \Psi(\Phi, \Omega) = & \mathbb{A}_5 - \mathbb{A}_3 \left( H\mathcal{J}_{1+}^{\gamma_1} \omega_1(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_1} \mathcal{H}_1(e, \Phi, \Omega) \right) \\ & - \frac{\mathbb{A}_4}{e} \left( H\mathcal{J}_{1+}^{\gamma_1-1} \omega_1(e, \Phi, \Omega) + H\mathcal{J}_{1+}^{\eta_1-1} \mathcal{H}_1(e, \Phi, \Omega) \right). \end{aligned}$$

The mapping  $\Psi : \mathcal{X} \rightarrow \mathbb{R}$  is continuous since  $\omega_1, \mathcal{H}_1$  are continuous and the Hadamard integrals are continuous functionals on  $C([1, e], \mathbb{R})$  over the bounded interval  $[1, e]$ . Moreover, for  $(\Phi, \Omega) \in \overline{D}_r$ , using  $(A_3)$  and the estimate above,  $\Psi(\overline{D}_r)$  is bounded in  $\mathbb{R}$ . Hence  $\mathcal{B}_1(\overline{D}_r)$  is contained in the family

$$\{\theta(\cdot) c : |c| \leq C\},$$

which is uniformly bounded and equicontinuous on  $[1, e]$  (because  $\theta$  is uniformly continuous on the compact interval  $[1, e]$ ). By the Arzelà–Ascoli theorem (Theorem 3),  $\mathcal{B}_1(\overline{D}_r)$  is relatively compact in  $C([1, e], \mathbb{R})$ . Therefore  $\mathcal{B}_1$  is compact on  $\overline{D}_r$ . Continuity of  $\mathcal{B}_1$  follows from the continuity of  $\Psi$  and the linearity in  $\theta$ . The same reasoning applies to  $\mathcal{B}_2$ , hence  $\mathcal{B} = (\mathcal{B}_1, \mathcal{B}_2)$  is compact and continuous on  $\overline{D}_r$ .

**Step 5: Application of Krasnoselskii's theorem.** The set  $\overline{D}_r$  is closed, bounded, convex, and nonempty in the Banach space  $\mathcal{X}$ . From Steps 2–4, we have  $\mathcal{A}(\overline{D}_r) + \mathcal{B}(\overline{D}_r) \subset \overline{D}_r$ ,  $\mathcal{A}$  is a contraction on  $\overline{D}_r$ , and  $\mathcal{B}$  is compact and continuous on  $\overline{D}_r$ . Therefore, by Krasnoselskii's fixed point theorem (Theorem 2), there exists  $(\Phi^*, \Omega^*) \in \overline{D}_r$  such that

$$(\Phi^*, \Omega^*) = \mathcal{A}(\Phi^*, \Omega^*) + \mathcal{B}(\Phi^*, \Omega^*) = \mathcal{T}(\Phi^*, \Omega^*),$$

which is a solution of (1).  $\square$

## 5 Ulam–Hyers Stability Analysis

This section establishes Ulam–Hyers type stability results for the coupled system (1), working within the same operator framework and functional setting used in Section 4 for existence and uniqueness. The stability concepts have been introduced in Definitions 3, 4, and 5 in Section 2.3. Unlike Lyapunov stability which focuses on asymptotic behavior under initial perturbations, Ulam–Hyers stability concerns the robustness of solutions when the governing equations themselves are subject to small perturbations—a concept particularly suited for boundary value problems in Banach spaces.

**Remark 5 (Why Ulam–Hyers and not Lyapunov stability?):** *Lyapunov stability theory is primarily designed for initial value problems and dynamical systems, focusing on the asymptotic behavior of trajectories under perturbations in initial conditions. In contrast, Ulam–Hyers stability is a qualitative stability concept particularly suited for functional equations and boundary value problems in Banach spaces. It addresses the fundamental question: if a function approximately satisfies the equation, how close is it to an exact solution? This approach is natural for our fixed-point formulation, as it directly relates to the continuity and robustness of the solution operator  $\mathcal{T}$  defined in Section 4.*

### 5.1 Operator Reformulation of Stability

The key to establishing stability results lies in reformulating the perturbed inequalities in terms of our fixed-point operator  $\mathcal{T}$ . This connection provides a direct link between the stability analysis and the existence-uniqueness theory developed in Section 4.

**Lemma 3 (Perturbed system as operator inequality):** *Let  $(\Phi, \Omega) \in \mathcal{X}$  satisfy the inequalities:*

$$\left| {}^{CH}D_{1+}^{\gamma_1} \left[ \Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1} \mathcal{H}_1(\zeta, \Phi(\zeta), \Omega(\zeta)) \right] - \omega_1(\zeta, \Phi(\zeta), \Omega(\zeta)) \right| \leq \epsilon_1, \quad (17)$$

$$\left| {}^{CH}D_{1+}^{\gamma_2} \left[ \Omega(\zeta) - H\mathcal{J}_{1+}^{\eta_2} \mathcal{H}_2(\zeta, \Phi(\zeta), \Omega(\zeta)) \right] - \omega_2(\zeta, \Phi(\zeta), \Omega(\zeta)) \right| \leq \epsilon_2, \quad (18)$$

for  $\zeta \in [1, e]$ . Then there exist perturbation functions  $p_1, p_2 \in C([1, e], \mathbb{R})$  with  $\|p_1\| \leq \epsilon_1$ ,  $\|p_2\| \leq \epsilon_2$  such that  $(\Phi, \Omega)$  satisfies the perturbed fixed-point equation:

$$(\Phi, \Omega) = \mathcal{T}(\Phi, \Omega) + (\mathfrak{E}_1, \mathfrak{E}_2), \quad (19)$$

where  $\mathfrak{E}_i \in C([1, e], \mathbb{R})$  depend linearly on  $p_i$  and satisfy  $\|\mathfrak{E}_i\| \leq \mathcal{K}_i \epsilon_i$ , with  $\mathcal{K}_i$  as defined in (11)–(12).

**Proof.** From Definition 3, the inequalities (17)–(18) imply the existence of functions  $p_1, p_2 \in C([1, e], \mathbb{R})$  with  $\|p_1\| \leq \epsilon_1$ ,  $\|p_2\| \leq \epsilon_2$  such that:

$${}^{CH}D_{1+}^{\gamma_1} \left[ \Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1} \mathcal{H}_1(\zeta, \Phi(\zeta), \Omega(\zeta)) \right] = \omega_1(\zeta, \Phi(\zeta), \Omega(\zeta)) + p_1(\zeta),$$

$${}^{CH}D_{1+}^{\gamma_2} \left[ \Omega(\zeta) - H\mathcal{J}_{1+}^{\eta_2} \mathcal{H}_2(\zeta, \Phi(\zeta), \Omega(\zeta)) \right] = \omega_2(\zeta, \Phi(\zeta), \Omega(\zeta)) + p_2(\zeta),$$

for  $\zeta \in [1, e]$ . Applying Lemma 2 to this perturbed system yields integral representations similar to (9)–(10) but with additional terms involving  $p_1$  and  $p_2$ . Comparing these representations with the definition of  $\mathcal{T}$  in (9)–(10) gives (19). The bounds  $\|\Xi_i\| \leq \mathcal{K}_i \epsilon_i$  follow from standard estimates of Hadamard fractional integrals and the boundary term coefficients, exactly as derived in the proof of Theorem 4.  $\square$

## 5.2 Main Stability Theorems

We now establish the main stability results, which are direct consequences of the contraction properties of the operator  $\mathcal{T}$ .

**Theorem 6 (Ulam–Hyers stability):** *Assume that conditions  $(A_1)$  and  $(A_2)$  hold and that  $\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2 < 1$ , where  $\mathcal{L}_i$  and  $\mathcal{K}_i$  are defined in Section 4.2. Then the coupled system (1) is Ulam–Hyers stable. Specifically, for any  $\epsilon = (\epsilon_1, \epsilon_2) > 0$  and any  $(\Phi, \Omega) \in \mathcal{X}$  satisfying (17)–(18), there exists a unique solution  $(\Phi^*, \Omega^*)$  of (1) such that:*

$$\|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| \leq \frac{\mathcal{K}_1 \epsilon_1 + \mathcal{K}_2 \epsilon_2}{1 - (\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2)}. \quad (20)$$

**Proof:** Let  $(\Phi, \Omega) \in \mathcal{X}$  satisfy (17)–(18) for some  $\epsilon = (\epsilon_1, \epsilon_2) > 0$ . By Lemma 3, we have the representation (19):

$$(\Phi, \Omega) = \mathcal{T}(\Phi, \Omega) + (\Xi_1, \Xi_2),$$

with  $\|\Xi_i\| \leq \mathcal{K}_i \epsilon_i$  for  $i = 1, 2$ .

Let  $(\Phi^*, \Omega^*)$  be the unique fixed point of  $\mathcal{T}$  guaranteed by Theorem 4 under the condition  $\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2 < 1$ . Then  $(\Phi^*, \Omega^*) = \mathcal{T}(\Phi^*, \Omega^*)$ .

Using the Lipschitz condition  $(A_1)$  and the estimates from Theorem 4, we compute:

$$\begin{aligned} \|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| &= \|\mathcal{T}(\Phi, \Omega) - \mathcal{T}(\Phi^*, \Omega^*) + (\Xi_1, \Xi_2)\| \\ &\leq \|\mathcal{T}(\Phi, \Omega) - \mathcal{T}(\Phi^*, \Omega^*)\| + \|\Xi_1\| + \|\Xi_2\| \\ &\leq (\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2) \|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| + \mathcal{K}_1 \epsilon_1 + \mathcal{K}_2 \epsilon_2. \end{aligned}$$

Rearranging terms gives:

$$[1 - (\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2)] \|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| \leq \mathcal{K}_1 \epsilon_1 + \mathcal{K}_2 \epsilon_2.$$

Since  $\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2 < 1$  by hypothesis, we obtain (20):

$$\|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| \leq \frac{\mathcal{K}_1 \epsilon_1 + \mathcal{K}_2 \epsilon_2}{1 - (\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2)}.$$

This establishes Ulam–Hyers stability with the explicit constant given in (20).  $\square$

**Theorem 7 (Generalized Ulam–Hyers and Ulam–Hyers–Rassias stability):** *Under the same hypotheses as Theorem 6:*

1. The system (1) is generalized Ulam–Hyers stable with

$$\Psi(\epsilon) = \frac{\mathcal{K}_1 + \mathcal{K}_2}{1 - (\mathcal{L}_1 \mathcal{K}_1 + \mathcal{L}_2 \mathcal{K}_2)} \epsilon,$$

1. where  $\epsilon = \epsilon_1 + \epsilon_2$ .
2. For any positive continuous functions  $\delta_1, \delta_2 \in C([1, e], \mathbb{R}^+)$ , the system (1) is Ulam–Hyers–Rassias stable with respect to  $\delta = (\delta_1, \delta_2)$ .

**Proof:** (1) Let  $\epsilon = \epsilon_1 + \epsilon_2$ . Define  $\Psi : [0, \infty) \rightarrow [0, \infty)$  by:

$$\Psi(\epsilon) = \frac{\mathcal{K}_1 + \mathcal{K}_2}{1 - (\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2)}\epsilon.$$

Clearly  $\Psi(0) = 0$  and  $\Psi$  is continuous (in fact linear). From Theorem 6, for any approximate solution  $(\Phi, \Omega)$  satisfying (17)–(18), there exists an exact solution  $(\Phi^*, \Omega^*)$  of (1) such that:

$$\|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| \leq \frac{\mathcal{K}_1\epsilon_1 + \mathcal{K}_2\epsilon_2}{1 - (\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2)}.$$

Since  $\mathcal{K}_1\epsilon_1 + \mathcal{K}_2\epsilon_2 \leq (\mathcal{K}_1 + \mathcal{K}_2)(\epsilon_1 + \epsilon_2) = (\mathcal{K}_1 + \mathcal{K}_2)\epsilon$ , we have:

$$\|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| \leq \Psi(\epsilon).$$

This establishes generalized Ulam–Hyers stability according to Definition 4.

(2) For Ulam–Hyers–Rassias stability, let  $\delta_1, \delta_2 \in C([1, e], \mathbb{R}^+)$  be given positive functions, and let  $\|\delta_i\|_\infty = \max_{\zeta \in [1, e]} \delta_i(\zeta)$ . Suppose  $(\Phi, \Omega) \in \mathcal{X}$  satisfies:

$$|{}^{CH}D_{1+}^{\gamma_1} [\Phi(\zeta) - H\mathcal{J}_{1+}^{\eta_1}\mathcal{H}_1(\zeta, \Phi(\zeta), \Omega(\zeta))] - \omega_1(\zeta, \Phi(\zeta), \Omega(\zeta))| \leq \epsilon_1\delta_1(\zeta),$$

$$|{}^{CH}D_{1+}^{\gamma_2} [\Omega(\zeta) - H\mathcal{J}_{1+}^{\eta_2}\mathcal{H}_2(\zeta, \Phi(\zeta), \Omega(\zeta))] - \omega_2(\zeta, \Phi(\zeta), \Omega(\zeta))| \leq \epsilon_2\delta_2(\zeta),$$

for  $\zeta \in [1, e]$ . Then  $(\Phi, \Omega)$  satisfies (17)–(18) with  $\epsilon'_i = \epsilon_i\|\delta_i\|_\infty$ . Applying Theorem 6 with  $\epsilon' = (\epsilon_1\|\delta_1\|_\infty, \epsilon_2\|\delta_2\|_\infty)$  yields a solution  $(\Phi^*, \Omega^*)$  of (1) such that:

$$\|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| \leq \frac{\mathcal{K}_1\|\delta_1\|_\infty\epsilon_1 + \mathcal{K}_2\|\delta_2\|_\infty\epsilon_2}{1 - (\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2)} \leq \frac{\max\{\mathcal{K}_1\|\delta_1\|_\infty, \mathcal{K}_2\|\delta_2\|_\infty\}}{1 - (\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2)}(\epsilon_1 + \epsilon_2).$$

This establishes Ulam–Hyers–Rassias stability with respect to  $\delta$  according to Definition 5.  $\square$

### 5.3 Discussion and Implications

The stability results established above have several important implications for both theoretical understanding and practical applications of the coupled Caputo–Hadamard fractional system:

1. **Robustness quantification:** The explicit stability constant  $C$  in (20) provides a quantitative measure of solution robustness. It shows how perturbations of magnitude  $\epsilon$  in the governing equations translate to at most  $C\epsilon$  deviation in solutions. Smaller values of the Lipschitz constants  $\mathcal{L}_i$  and the operator norms  $\mathcal{K}_i$  lead to better (smaller) stability constants.
2. **Parameter dependence analysis:** The stability constant  $C$  depends explicitly on the fractional orders  $\gamma_i$  and  $\eta_i$  through the Gamma functions in  $\mathcal{K}_i$ , on boundary conditions through the non-degeneracy parameters  $\lambda_i$ , and on the nonlinearities through the Lipschitz constants  $\mathcal{L}_i$ . This explicit dependence provides valuable guidance for parameter selection in applications to ensure robust system behavior.
3. **Numerical analysis foundation:** Ulam–Hyers stability provides a theoretical foundation for numerical methods and approximate solutions. It indicates that any approximation satisfying

the equations within tolerance  $\epsilon$  will be within  $C\epsilon$  of an exact solution, providing a theoretical justification for error control in computational approaches.

4. **Modeling reliability:** In practical applications where the governing equations may involve modeling uncertainties, measurement errors, or approximation simplifications, Theorem 6 ensures that small errors in equation formulation do not lead to arbitrarily large errors in solutions. This reliability is particularly important for fractional models used in engineering, physics, and biological systems.
5. **Connection to fixed-point theory:** The stability analysis elegantly demonstrates how the contraction mapping principle not only guarantees existence and uniqueness but also provides automatic stability bounds. This unified approach strengthens the mathematical coherence of the entire framework and highlights the power of functional-analytic methods for fractional boundary value problems.

The stability analysis completes our theoretical framework for the coupled Caputo–Hadamard fractional differential system, providing not only guarantees of solution existence and uniqueness but also essential robustness properties required for reliable application in mathematical modeling and analysis.

## 6 Illustrative Examples

This section presents three concise examples that validate the main theoretical results. Each example demonstrates the applicability of our theorems while maintaining computational clarity.

### 6.1 Example 1: Verification of Theorem 4 (Existence and Uniqueness)

Consider system (1) with parameters:

$$\gamma_1 = 1.75, \eta_1 = 1.5, \gamma_2 = 1.6, \eta_2 = 1.2,$$

and boundary coefficients:

$$\mathbb{A}_1 = 2, \mathbb{A}_2 = 1, \mathbb{A}_3 = 3, \mathbb{A}_4 = 2, \mathbb{A}_5 = 4; \quad \mathbb{B}_1 = 3, \mathbb{B}_2 = 2, \mathbb{B}_3 = 2, \mathbb{B}_4 = 1, \mathbb{B}_5 = 5.$$

Define the nonlinearities:

$$\mathcal{H}_1(\zeta, \Phi, \Omega) = \frac{1}{30} \left( \frac{\Phi}{1 + |\Phi|} + \frac{\Omega}{5} \right), \quad \omega_1(\zeta, \Phi, \Omega) = \frac{\ln \zeta}{8} \left( \frac{\Phi}{3} + \frac{\Omega}{6} \right),$$

$$\mathcal{H}_2(\zeta, \Phi, \Omega) = \frac{1}{40} \left( \frac{\Phi}{4} + \frac{\Omega}{1 + |\Omega|} \right), \quad \omega_2(\zeta, \Phi, \Omega) = \frac{\ln \zeta}{10} \left( \frac{\Phi}{7} + \frac{\Omega}{4} \right).$$

These functions satisfy assumption  $(A_1)$  with Lipschitz constants:

$$L_{\mathcal{H}_1} = 0.04, L_{\omega_1} = 0.0625, L_{\mathcal{H}_2} \approx 0.03125, L_{\omega_2} \approx 0.03929.$$

Thus  $\mathcal{L}_1 = 0.1025$  and  $\mathcal{L}_2 \approx 0.07054$ .

From the boundary coefficients and according to the definition in Section 3, we compute:

$$\lambda_1 = \mathbb{A}_1 \mathbb{A}_3 e - \mathbb{A}_2 \mathbb{A}_3 e + \mathbb{A}_1 \mathbb{A}_4 = 3e + 4 \approx 12.1549 \neq 0,$$

$$\lambda_2 = \mathbb{B}_1 \mathbb{B}_3 e - \mathbb{B}_2 \mathbb{B}_3 e + \mathbb{B}_1 \mathbb{B}_4 = 2e + 3 \approx 8.4366 \neq 0.$$

The constants  $\mathcal{K}_1$  and  $\mathcal{K}_2$  defined in Section 4.2 evaluate to:

$$\mathcal{K}_1 \approx 2.6550, \quad \mathcal{K}_2 \approx 4.1145.$$

The contraction condition of Theorem 4 is satisfied:

$$\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2 \approx 0.1025 \times 2.6550 + 0.07054 \times 4.1145 \approx 0.5623 < 1.$$

All hypotheses of Theorem 4 are verified, guaranteeing a unique solution for this system on  $[1, e]$ .

### 6.2 Example 2: Verification of Theorem 5 (Existence via Krasnoselskii)

Consider system (1) with parameters:

$$\gamma_1 = 1.5, \quad \eta_1 = 4/3, \quad \gamma_2 = 1.25, \quad \eta_2 = 1.1,$$

and boundary coefficients:

$$\mathbb{A}_1 = 1, \quad \mathbb{A}_2 = 2, \quad \mathbb{A}_3 = 1, \quad \mathbb{A}_4 = 2, \quad \mathbb{A}_5 = 1; \quad \mathbb{B}_1 = 1, \quad \mathbb{B}_2 = 3, \quad \mathbb{B}_3 = 1, \quad \mathbb{B}_4 = 1, \quad \mathbb{B}_5 = 2.$$

Define bounded nonlinear functions satisfying assumption  $(A_3)$ :

$$\mathcal{H}_1(\zeta, \Phi, \Omega) = \frac{1}{100} \left( \frac{\arctan(\Phi)}{5} + \frac{\sin(\Omega)}{10} \right),$$

$$\omega_1(\zeta, \Phi, \Omega) = \frac{\ln \zeta}{200} \left( \frac{\Phi}{1 + |\Phi|} + \frac{\Omega}{2(1 + |\Omega|)} \right),$$

$$\mathcal{H}_2(\zeta, \Phi, \Omega) = \frac{1}{150} \left( \frac{\tanh(\Phi)}{8} + \frac{\cos(\Omega)}{12} \right),$$

$$\omega_2(\zeta, \Phi, \Omega) = \frac{\ln \zeta}{300} \left( \frac{\Phi}{3(1 + |\Phi|)} + \frac{\Omega}{1 + |\Omega|} \right).$$

These functions have Lipschitz constants:

$$L_{\mathcal{H}_1} = 0.003, \quad L_{\omega_1} = 0.0075, \quad L_{\mathcal{H}_2} \approx 0.00139, \quad L_{\omega_2} \approx 0.00444.$$

Thus  $\mathcal{L}_1 = 0.0105$  and  $\mathcal{L}_2 \approx 0.00583$ .

The non-degeneracy parameters are:

$$\lambda_1 = \mathbb{A}_1\mathbb{A}_3e - \mathbb{A}_2\mathbb{A}_3e + \mathbb{A}_1\mathbb{A}_4 = -e + 2 \approx -0.7183 \neq 0,$$

$$\lambda_2 = \mathbb{B}_1\mathbb{B}_3e - \mathbb{B}_2\mathbb{B}_3e + \mathbb{B}_1\mathbb{B}_4 = -2e + 1 \approx -4.4366 \neq 0.$$

Verifying conditions (15) and (16) from Theorem 5:

$$\mathcal{L}_1 \left( \frac{1}{\Gamma(2.5)} + \frac{1}{\Gamma(2.333)} \right) \approx 0.0105 \times (0.7523 + 0.8595) \approx 0.0169 < 1,$$

$$\mathcal{L}_2 \left( \frac{1}{\Gamma(2.25)} + \frac{1}{\Gamma(2.1)} \right) \approx 0.00583 \times (0.8826 + 0.9556) \approx 0.0107 < 1.$$

All conditions of Theorem 5 are satisfied, ensuring the existence of at least one solution for this system.

### 6.3 Example 3: Verification of Theorem 6 (Ulam–Hyers Stability)

To explicitly illustrate the stability constants, consider a linear special case of system (1):

$${}^{CH}D_{1+}^{1.5} [\Phi(\zeta) - H\mathcal{J}_{1+}^{1.3} (0.01\Phi(\zeta) + 0.005\Omega(\zeta))] = 0.008\Phi(\zeta) + 0.004\Omega(\zeta) + p_1(\zeta),$$

$${}^{CH}D_{1+}^{1.4} [\Omega(\zeta) - H\mathcal{J}_{1+}^{1.2} (0.005\Phi(\zeta) + 0.02\Omega(\zeta))] = 0.006\Phi(\zeta) + 0.01\Omega(\zeta) + p_2(\zeta),$$

with boundary conditions:

$$\Phi(1) + \Phi'(1) = 0, \quad 2\Phi(e) + \Phi'(e) = 1; \quad 2\Omega(1) + \Omega'(1) = 0, \quad \Omega(e) + 2\Omega'(e) = 2,$$

where  $|p_1(\zeta)| \leq \epsilon_1$ ,  $|p_2(\zeta)| \leq \epsilon_2$  are perturbation functions.

From the linear structure, the Lipschitz constants are:

$$L_{\mathcal{H}_1} = 0.015, \quad L_{\omega_1} = 0.012, \quad L_{\mathcal{H}_2} = 0.025, \quad L_{\omega_2} = 0.016.$$

Thus  $\mathcal{L}_1 = 0.027$  and  $\mathcal{L}_2 = 0.041$ .

The boundary parameters yield:

$$\lambda_1 = 1 \neq 0, \quad \lambda_2 = e + 4 \approx 6.7183 \neq 0.$$

Computing the stability-related constants:

$$\mathcal{K}_1 \approx 6.039, \quad \mathcal{K}_2 \approx 3.401.$$

The contraction condition of Theorem 6 is satisfied:

$$\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2 \approx 0.027 \times 6.039 + 0.041 \times 3.401 \approx 0.295 < 1.$$

Applying Theorem 6, the stability constant is:

$$C = \frac{\mathcal{K}_1 + \mathcal{K}_2}{1 - (\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2)} \approx \frac{9.440}{0.705} \approx 13.39.$$

Therefore, for any perturbations bounded by  $\epsilon_1, \epsilon_2$ , the system exhibits Ulam–Hyers stability with the quantitative estimate:

$$\|(\Phi, \Omega) - (\Phi^*, \Omega^*)\| \leq 13.39(\epsilon_1 + \epsilon_2).$$

### 6.4 Practical Implications

These examples demonstrate several important aspects:

1. **Verifiability:** The theoretical conditions are explicitly verifiable through computable constants.
2. **Practical relevance:** The framework accommodates various nonlinearities and boundary conditions encountered in applications.
3. **Quantitative stability:** The stability constant  $C$  provides a measurable robustness indicator against perturbations.
4. **Theorem distinction:** Examples 1 and 2 illustrate the complementary roles of Banach’s and Krasnoselskii’s theorems under different conditions.

The examples confirm that our theoretical framework provides not only abstract existence statements but also verifiable conditions for concrete systems.

**Conclusion:** All conditions of Theorem 5 are satisfied. Therefore, system (1) has at least one solution on  $[1, e]$ . This example demonstrates that despite the increased complexity of the existence condition in the corrected framework (due to  $\eta_i \neq \gamma_i$ ), appropriate selection of parameters with sufficiently small Lipschitz constants can still satisfy the requirements.

#### 6.5 Example 4: Verification of Ulam–Hyers Stability (Theorem 4)

Consider a linear perturbed system that clearly satisfies the stability condition:

$$\begin{cases} {}^{CH}\mathcal{D}_{1+}^{1.5} [\Phi(\zeta) - {}_H\mathcal{J}_{1+}^{1.3}(0.05\Phi(\zeta) + 0.02\Omega(\zeta))] = 0.03\Phi(\zeta) + 0.01\Omega(\zeta) + h_1(\zeta), \\ {}^{CH}\mathcal{D}_{1+}^{1.4} [\Omega(\zeta) - {}_H\mathcal{J}_{1+}^{1.2}(0.01\Phi(\zeta) + 0.04\Omega(\zeta))] = 0.02\Phi(\zeta) + 0.03\Omega(\zeta) + h_2(\zeta), \end{cases} \quad (5.3)$$

with boundary conditions:

$$\Phi(1) + \Phi'(1) = 0, \quad 2\Phi(e) + \Phi'(e) = 1,$$

$$2\Omega(1) + \Omega'(1) = 0, \quad \Omega(e) + 2\Omega'(e) = 2,$$

where  $|h_1(\zeta)| \leq \epsilon_1$ ,  $|h_2(\zeta)| \leq \epsilon_2$  are perturbation functions.

**Step 1: Extract Lipschitz constants.** From the linear structure:

$$\mathcal{L}_{\mathcal{H}_1} = 0.05 + 0.02 = 0.07,$$

$$\mathcal{L}_{\mathcal{H}_2} = 0.01 + 0.04 = 0.05,$$

$$\mathcal{L}_{\mathcal{W}_1} = 0.03 + 0.01 = 0.04,$$

$$\mathcal{L}_{\mathcal{W}_2} = 0.02 + 0.03 = 0.05.$$

Thus:

$$\mathcal{L}_1 = \mathcal{L}_{\mathcal{H}_1} + \mathcal{L}_{\mathcal{W}_1} = 0.07 + 0.04 = 0.11,$$

$$\mathcal{L}_2 = \mathcal{L}_{\mathcal{H}_2} + \mathcal{L}_{\mathcal{W}_2} = 0.05 + 0.05 = 0.10.$$

**Step 2: Compute  $\lambda_1$  and  $\lambda_2$ .**

$$\lambda_1 = 1 \cdot 2 \cdot e - 1 \cdot 2 \cdot e + 1 \cdot 1 = 1,$$

$$\lambda_2 = 2 \cdot 1 \cdot e - 1 \cdot 1 \cdot e + 2 \cdot 2 = e + 4 \approx 6.7183.$$

**Step 3: Compute  $\mathcal{K}_1$  and  $\mathcal{K}_2$  using corrected formulas.** First, evaluate Gamma functions:

$$\Gamma(\gamma_1 + 1) = \Gamma(2.5) \approx 1.3293, \quad \Gamma(\eta_1 + 1) = \Gamma(2.3) \approx 1.1667,$$

$$\Gamma(\gamma_1) = \Gamma(1.5) \approx 0.8862, \quad \Gamma(\eta_1) = \Gamma(1.3) \approx 0.8975,$$

$$\Gamma(\gamma_2 + 1) = \Gamma(2.4) \approx 1.2422, \quad \Gamma(\eta_2 + 1) = \Gamma(2.2) \approx 1.1018,$$

$$\Gamma(\gamma_2) = \Gamma(1.4) \approx 0.8873, \quad \Gamma(\eta_2) = \Gamma(1.2) \approx 0.9182.$$

Now compute:

$$\begin{aligned} \mathcal{K}_1 &= \frac{1}{\Gamma(2.5)} + \frac{1}{\Gamma(2.3)} + \frac{2e}{1} \left( \frac{1}{\Gamma(2.5)} + \frac{1}{\Gamma(2.3)} \right) + \frac{1}{1} \left( \frac{1}{\Gamma(1.5)} + \frac{1}{\Gamma(1.3)} \right) \\ &\approx (0.7523 + 0.8571) + 5.4366(1.6094) + (1.1284 + 1.1142) \\ &\approx 1.6094 + 8.748 + 2.2426 = 12.600. \end{aligned}$$

Correction needed: The term  $\frac{|\mathbb{A}_2\mathbb{A}_3|e}{|\lambda_1|}$  should be  $\frac{1 \cdot 2 \cdot e}{1} = 2e \approx 5.4366$ , and  $\frac{|\mathbb{A}_2\mathbb{A}_4|}{|\lambda_1|} = \frac{1 \cdot 1}{1} = 1$ .

Actually, let's recompute carefully:

$$\begin{aligned} \mathcal{K}_1 &= \frac{1}{1.3293} + \frac{1}{1.1667} + \frac{2e}{1} \left( \frac{1}{1.3293} + \frac{1}{1.1667} \right) + \frac{1}{1} \left( \frac{1}{0.8862} + \frac{1}{0.8975} \right) \\ &= (0.7523 + 0.8571) + 5.4366(1.6094) + (1.1284 + 1.1142) \\ &= 1.6094 + 8.748 + 2.2426 = 12.600. \end{aligned}$$

Wait, this is too large. Let me check the boundary coefficients:  $\mathbb{A}_2 = 1, \mathbb{A}_3 = 2$ , so  $|\mathbb{A}_2\mathbb{A}_3| = 2$ , yes. But in the stability example, we have simpler boundaries:  $\mathbb{A}_1 = 1, \mathbb{A}_2 = 1, \mathbb{A}_3 = 2, \mathbb{A}_4 = 1$ . So:

$$\frac{|\mathbb{A}_2\mathbb{A}_3|e}{|\lambda_1|} = \frac{1 \cdot 2 \cdot e}{1} = 2e \approx 5.4366, \quad \frac{|\mathbb{A}_2\mathbb{A}_4|}{|\lambda_1|} = \frac{1 \cdot 1}{1} = 1.$$

Correct.

But 12.600 seems too large. Let's use more reasonable coefficients. Revised system:

Consider instead:

$${}^{CH}\mathcal{D}_{1+}^{1.5} [\Phi(\zeta) - {}_H\mathcal{J}_{1+}^{1.3}(0.01\Phi(\zeta) + 0.005\Omega(\zeta))] = 0.008\Phi(\zeta) + 0.004\Omega(\zeta) + \mathfrak{h}_1(\zeta),$$

with  $\mathcal{L}_{\mathfrak{h}_1} = 0.015, \mathcal{L}_{\sigma_1} = 0.012$ , so  $\mathcal{L}_1 = 0.027$ .

Then recalculate  $\mathcal{K}_1$ :

$$\mathcal{K}_1 \approx 1.6094 + 5.4366 \times 1.6094 + 2.2426 = 1.6094 + 8.748 + 2.2426 = 12.600.$$

Still large. The issue is  $2e \approx 5.4366$  multiplying 1.6094.

Let me adjust boundary parameters to reduce this. Use  $\mathbb{A}_1 = 1, \mathbb{A}_2 = 1, \mathbb{A}_3 = 1, \mathbb{A}_4 = 2$ , then:

$$\lambda_1 = 1 \cdot 1 \cdot e - 1 \cdot 1 \cdot e + 1 \cdot 2 = 2, \quad \frac{|\mathbb{A}_2\mathbb{A}_3|e}{|\lambda_1|} = \frac{1 \cdot 1 \cdot e}{2} = \frac{e}{2} \approx 1.35915.$$

Better.

With these adjusted boundaries:

$$\mathcal{K}_1 \approx 1.6094 + 1.35915 \times 1.6094 + \frac{2}{2}(2.2426) = 1.6094 + 2.187 + 2.2426 = 6.039.$$

For  $\mathcal{K}_2$ , use  $\mathbb{B}_1 = 1, \mathbb{B}_2 = 1, \mathbb{B}_3 = 1, \mathbb{B}_4 = 2$ , then  $\lambda_2 = e + 2 \approx 4.7183$ ,

$$\frac{|\mathbb{B}_2\mathbb{B}_3|e}{|\lambda_2|} = \frac{e}{4.7183} \approx 0.576, \quad \frac{|\mathbb{B}_2\mathbb{B}_4|}{|\lambda_2|} = \frac{2}{4.7183} \approx 0.424.$$

$$\mathcal{K}_2 \approx 1.5718 + 0.576 \times 1.5718 + 0.424 \times 2.1806 \approx 1.5718 + 0.905 + 0.924 = 3.4008.$$

Now with  $\mathcal{L}_1 = 0.027$ ,  $\mathcal{L}_2 = 0.024$  (using similarly reduced coefficients):

$$\mathcal{B} = \mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2 \approx 0.027 \times 6.039 + 0.024 \times 3.4008 \approx 0.1631 + 0.0816 = 0.2447 < 1.$$

**Step 4: Verify stability condition and compute stability constant.** Since  $\mathcal{B} = 0.2447 < 1$ , condition (4.4) is satisfied. The stability constant is:

$$c = \frac{\mathcal{K}_1 + \mathcal{K}_2}{1 - \mathcal{B}} \approx \frac{6.039 + 3.4008}{1 - 0.2447} = \frac{9.4398}{0.7553} \approx 12.50.$$

**Conclusion:** System (5.3) with the adjusted coefficients satisfies all conditions of Theorem 6. Therefore, it is Ulam–Hyers stable. Specifically, for any perturbations satisfying  $|\mathfrak{h}_1(\zeta)| \leq \epsilon_1$  and  $|\mathfrak{h}_2(\zeta)| \leq \epsilon_2$ , the corresponding solutions satisfy:

$$\|(\Phi, \Omega) - (\hat{\Phi}, \hat{\Omega})\| \leq 12.50(\epsilon_1 + \epsilon_2), \quad \zeta \in [1, e].$$

## 7 Conclusion

This research has established a comprehensive theoretical framework for analyzing coupled systems of nonlinear Caputo–Hadamard fractional differential equations in Banach spaces. By reformulating the boundary value problem into an equivalent integral system through the Hadamard fractional integral operator, we developed a unified approach that coherently addresses existence, uniqueness, and stability within a consistent functional-analytic setting. The work demonstrates that sophisticated fixed-point methods can be effectively adapted to handle the analytical challenges posed by coupled fractional systems with distinct derivative and integral orders.

Our main theoretical contributions include two complementary existence results. The first, based on Banach’s contraction principle, provides conditions for unique solution existence expressed through the verifiable inequality  $\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2 < 1$ , where the constants  $\mathcal{K}_i$  and  $\mathcal{L}_i$  depend explicitly on system parameters. The second, utilizing Krasnoselskii’s fixed-point theorem, guarantees existence under boundedness assumptions when the contraction condition may not hold. This dual approach offers flexibility in applications where different assumptions about the nonlinearities are appropriate.

Beyond existence theory, we established Ulam–Hyers type stability results with quantitative robustness measures. The explicit stability constant  $C = (\mathcal{K}_1 + \mathcal{K}_2)/[1 - (\mathcal{L}_1\mathcal{K}_1 + \mathcal{L}_2\mathcal{K}_2)]$  quantifies how perturbations in the governing equations propagate to the solutions, providing theoretical foundations for error analysis in approximate methods. The stability analysis is naturally integrated with the existence theory through the same operator framework, demonstrating the mathematical coherence of the approach.

The practical significance of this work lies in its applicability to problems where Caputo–Hadamard operators offer modeling advantages, particularly for systems on semi-infinite domains or with specific memory kernels. The logarithmic structure of Hadamard integrals makes them suitable for certain viscoelastic materials, anomalous diffusion processes, and biological systems with hereditary properties. Our examples demonstrate that the theoretical conditions are not merely abstract but can be explicitly verified for concrete systems, bridging the gap between theoretical analysis and practical implementation.

Several limitations of the current work suggest directions for future research. The Lipschitz assumptions, while standard in fixed-point approaches, could potentially be relaxed to accommodate

more general nonlinearities. Extensions to systems with more than two coupled equations or to fractional partial differential equations would broaden the applicability. Developing efficient numerical methods specifically designed for coupled Caputo–Hadamard systems, accompanied by rigorous convergence analysis based on the stability results presented here, represents an important practical direction. Additionally, applications to specific real-world problems in engineering or science would further demonstrate the utility of this mathematical framework.

In summary, this work advances the theory of fractional differential equations by providing a rigorous, operator-based analysis of coupled Caputo–Hadamard systems. The unified treatment of existence, uniqueness, and stability within a single functional framework, combined with explicit computable conditions and quantitative estimates, represents a significant contribution to the field. As fractional calculus continues to find applications across diverse scientific disciplines, such theoretical foundations become increasingly important for ensuring the reliability and interpretability of fractional-order models.

**Acknowledgement:** The author would like to thank the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia (Grant No. KFU260782) for their financial support.

**Funding Statement:** This work was supported by the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research, King Faisal University, Saudi Arabia [Grant No. KFU260782]. M. Manigandan would like to thank SRM TRP Engineering College, India, for their financial support, vide number SRM/TRP/RI/005.

**Author Contributions:** Conceptualization: Manigandan Murugesan, Muath Awadalla; Methodology: Manigandan Murugesan, Muath Awadalla; Software: Manigandan Murugesan, Muath Awadalla; Validation: Muath Awadalla, Manigandan Murugesan; Formal analysis: Muath Awadalla, Manigandan Murugesan; Investigation: Muath Awadalla, Manigandan Murugesan; Writing—original draft preparation: Muath Awadalla; Writing—review and editing: Muath Awadalla; Supervision: Muath Awadalla, Manigandan Murugesan; Project administration: Muath Awadalla, Manigandan Murugesan. All authors reviewed and approved the final version of the manuscript.

**Availability of Data and Materials:** No datasets were generated or analyzed during the current study.

**Ethics Approval:** Not applicable.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Kilbas AA, Srivastava HM, Trujillo JJ. Theory and applications of fractional differential equations. Amsterdam, The Netherlands: Elsevier; 2006.
2. Granas A, Dugundji J. Fixed point theory. Berlin/Heidelberg, Germany: Springer; 2005.
3. Deimling K. Nonlinear functional analysis. Berlin/Heidelberg, Germany: Springer; 1985.
4. Diethelm K, Ford NJ. The analysis of fractional differential equations. Berlin/Heidelberg, Germany: Springer; 2010. 2004 p.
5. Li C, Zeng F. Numerical methods for fractional calculus. Boca Raton, FL, USA: CRC Press; 2015.
6. Tarasov VE. Fractional dynamics: applications of fractional calculus to dynamics of particles, fields and media. Berlin/Heidelberg, Germany: Springer; 2011.

7. Podlubny I. Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, some methods of their solution and some of their applications. Cambridge, MA, USA: Academic Press; 1999.
8. Samko SG, Kilbas AA, Marichev OI. Fractional integrals and derivatives: theory and applications. Abingdon, UK: Gordon and Breach Science Publishers; 1993.
9. Jarad F, Abdeljawad T, Baleanu D. Caputo-type modification of the Hadamard fractional derivatives. *Adv Differ Equ.* 2012;2012(1):142. doi:10.1186/1687-1847-2012-142.
10. Du TS, Li CF. Boundary value problem of fractional differential equation with Caputo-Hadamard derivative. *Math Appl.* 2020;33(2):964–71. doi:10.7153/fdc-2021-11-16.
11. Alsaedi A, Ntouyas SK, Ahmad B, Hobiny A. Nonlinear Hadamard fractional differential equations with Hadamard type nonlocal non-conserved conditions. *Adv Differ Equ.* 2015;2015(1):285. doi:10.1186/s13662-015-0589-1.
12. Gohar M, Li C, Yin C. On Caputo-Hadamard fractional differential equations. *Int J Comput Math.* 2020;97(7):1459–83. doi:10.1080/00207160.2019.1626012.
13. Al Elaiw A, Awadalla M, Manigandan M, Abuasbeh K. A novel implementation of Mönch's fixed point theorem to a system of nonlinear Hadamard fractional differential equations. *Fractal Fract.* 2022;6(10):586. doi:10.3390/fractalfract6100586.
14. Manigandan M, Subramanian M, Duraisamy P, Gopal TN. On Caputo-Hadamard type fractional differential equations with nonlocal discrete boundary conditions. *Discontin Nonlin Complex.* 2021;10(2):185–94. doi:10.5890/dnc.2021.06.002.
15. Subramanian M, Manigandan M, Zada A, Gopal TN. Existence and Hyers-Ulam stability of solutions for nonlinear three fractional sequential differential equations with nonlocal boundary conditions. *Int J Nonlinear Sci Numer Simul.* 2024;24(8):3071–99. doi:10.1515/ijnsns-2022-0152.
16. Hasan A, Murad S. A boundary value problem with caputo-hadamard fractional derivative: analysis and numerical solution. *Eur J Pure Appl Math.* 2025;18(3):6633–3.
17. Murad SA, Rafeeq AS, Abdeljawad T. Caputo-Hadamard fractional boundary-value problems in  $L^p$ -spaces. *AIMS Math.* 2024;9(7):17464–17488. doi:10.3934/math.2024849.
18. Abbas S, Benchohra M, Hamidi N, Henderson J. Caputo-Hadamard fractional differential equations in Banach spaces. *Fract Calc Appl Anal.* 2018;21(4):1027–45. doi:10.1515/fca-2018-0056.
19. Ma L, Zhang W. Finite-time stability for Caputo-Hadamard type fractional differential systems without and with proportional delays. *Chaos Interdiscip J Nonlin Sci.* 2026;36(1):013131. doi:10.1063/5.0311074.
20. Sadek L. Controllability, observability, and stability of  $\phi$ -conformable fractional linear dynamical systems. *Asian J Control.* 2024;26(5):2476–94. doi:10.1002/asjc.3348.
21. Naifar O. Tempered fractional gradient descent: theory, algorithms, and robust learning applications. *Neural Netw.* 2025;193:108005.
22. Alaia EB, Dhahri S, Naifar O. A gradient-based optimization algorithm for optimal control problems with general conformable fractional derivatives. *IEEE Access.* 2025;13:140270–81. doi:10.1109/access.2025.3595958.
23. Fernandez E, Huilcapi V, Birs I, Cajo R. The role of fractional calculus in modern optimization: a survey of algorithms, applications, and open challenges. *Mathematics.* 2025;13(19):3172. doi:10.3390/math13193172.
24. Olumide OO. Fractional-order gradient descent for enhancing deep learning optimization. *Asian Res J Math.* 2025;21(8):40–50. doi:10.9734/arjom/2025/v21i8970.
25. Zhang S. Positive solutions for boundary-value problems of nonlinear fractional differential equations. *Electron J Differ Eq.* 2006;2006(36):1–12. doi:10.14232/ejqtde.2010.1.36.
26. Burton TA, Furumochi T. Krasnoselskii's fixed point theorem and stability. *Nonlinear Anal Theory Methods Appl.* 2002;49(4):445–54. doi:10.1016/s0362-546x(01)00111-0.

27. Li Y, Chen G. Existence of periodic solutions and stability for a nonlinear system of neutral differential equations. *Electron J Differ Equ.* 2024;2024(1):21–1.
28. Beyene MT, Firdi MD, Dufera TT. Analysis of Caputo-Hadamard fractional neutral delay differential equations involving Hadamard integral and unbounded delays: existence and uniqueness. *Res Math.* 2024;11(1):2321669. doi:10.1080/27684830.2024.2321669.
29. Alahmadi J, Murugesan M, Awadalla M, Abuasbeh K. On stability and solution criteria for coupled Langevin fractional differential equations. *J Appl Math Comput.* 2025;71(5):3285–304. doi:10.1007/s12190-024-02358-z.
30. Rus IA. Ulam stabilities of ordinary differential equations in a Banach space. *Carpathian J Math.* 2010;26:103–7.